

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 June 2002 (20.06.2002)

PCT

(10) International Publication Number
WO 02/48189 A2

(51) International Patent Classification⁷: C07K 14/47

(74) Agent: HOFMAN-BANG ZACCO A/S; Hans
Bekkevolds Allé 7, DK-2900 Hellerup (DK).

(21) International Application Number: PCT/DK01/00825

(22) International Filing Date:
13 December 2001 (13.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2000 01872 13 December 2000 (13.12.2000) DK
60/272,098 28 February 2001 (28.02.2001) US

(71) Applicant (*for all designated States except US*): BOREAN
PHARMA A/S [DK/DK]; Science Park Aarhus, Gustav
Wieds Vej 10, DK-8000 Aarhus C (DK).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ETZERODT,
Michael [DK/DK]; Teglhøj 18, DK-8382 Hinnerup
(DK). HOLTET, Thor, Las [DK/DK]; Roskildevej
104, DK-3600 Frederikssund (DK). GRAVERSEN,
Niels, Jonas, Heilskov [DK/DK]; Brendstrupvej 58,
DK-8200 Århus N (DK). THØGERSEN, Hans, Chris-
tian [DK/DK]; Ristrupvej 41, DK-8381 Mundelstrup
(DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: COMBINATORIAL LIBRARIES OF PROTEINS HAVING THE SCAFFOLD STRUCTURE OF C-TYPE LECTIN-
LIKE DOMAINS

(57) Abstract: A novel family of protein libraries comprising CTLDS (C-type Lectin-Like Domains) in which internal polypeptide loop-regions lining the ligand binding sites in CTLDS have been replaced with ensembles of completely or partially randomised polypeptide segments. Tetranectin CTLDS were chosen as framework for the preferred embodiment of the invention; and versatile phagemid vectors useful in the generation and manipulation of human and murine tetranectin CTLD libraries are disclosed as part of this invention. Tetranectin CTLDS in monomeric as well as in trimeric form are efficiently displayed as gene III fusions in fully functional form by the recombinant fd phage display vector. CTLD derivatives with affinity for new ligands may readily be isolated from libraries of vectors displaying CTLDS, in which loop-regions have been randomised, using one or more rounds of enrichment by screening or selection followed by amplification of the enriched subpopulation in each round. The efficiency with which protein products containing CTLDS with new binding properties can be produced, e.g. by bacterial expression in *in vitro* refolding, in mono-, tri-, or multimeric formats provides important advantages in terms of simplicity, cost and efficiency of generation, production and diagnostic or therapeutic applications in comparison to recombinant antibody derivatives.

WO 02/48189 A2

Combinatorial libraries of proteins having the scaffold
structure of C-type lectin-like domains

FIELD OF THE INVENTION

5 This invention describes a system which relates to the generation of randomised libraries of ligand-binding protein units derived from proteins containing the so-called C-type lectin like domain (CTLD) of which the carbohydrate recognition domain (CRD) of C-type lectins represents one example
10 of a family of this protein domain.

BACKGROUND OF THE INVENTION

The C-type lectin-like domain (CTLD) is a protein domain family which has been identified in a number of proteins isolated from many animal species (reviewed in Drickamer
15 and Taylor (1993) and Drickamer (1999)). Initially, the CTLD domain was identified as a domain common to the so-called C-type lectins (calcium-dependent carbohydrate binding proteins) and named "Carbohydrate Recognition Domain" ("CRD"). More recently, it has become evident that this domain
20 is shared among many eukaryotic proteins, of which several do not bind sugar moieties, and hence, the canonical domain has been named as CTLD.

CTLDs have been reported to bind a wide diversity of compounds, including carbohydrates, lipids, proteins, and even
25 ice [Aspberg et al. (1997), Bettler et al. (1992), Ewart et al. (1998), Graversen et al. (1998), Mizumo et al. (1997), Sano et al. (1998), and Tormo et al. (1999)]. Only one copy of the CTLD is present in some proteins, whereas other proteins contain from two to multiple copies of the domain. In
30 the physiologically functional unit multiplicity in the number of CTLDs is often achieved by assembling single copy protein protomers into larger structures.

The CTLD consists of approximately 120 amino acid residues and, characteristically, contains two or three intra-chain disulfide bridges. Although the similarity at the amino acid sequence level between CTLDs from different proteins is relatively low, the 3D-structures of a number of CTLDs have been found to be highly conserved, with the structural variability essentially confined to a so-called loop-region, often defined by up to five loops. Several CTLDs contain either one or two binding sites for calcium and most of the side chains which interact with calcium are located in the loop-region.

On the basis of CTLDs for which 3D structural information is available, it has been inferred that the canonical CTLD is structurally characterised by seven main secondary-structure elements (i.e. five β -strands and two α -helices) sequentially appearing in the order $\beta 1$; $\alpha 1$; $\alpha 2$; $\beta 2$; $\beta 3$; $\beta 4$; and $\beta 5$ (Fig. 1, and references given therein). In all CTLDs, for which 3D structures have been determined, the β -strands are arranged in two anti-parallel β -sheets, one composed of $\beta 1$ and $\beta 5$, the other composed of $\beta 2$, $\beta 3$ and $\beta 4$. An additional β -strand, $\beta 0$, often precedes $\beta 1$ in the sequence and, where present, forms an additional strand integrating with the $\beta 1$, $\beta 5$ -sheet. Further, two disulfide bridges, one connecting $\alpha 1$ and $\beta 5$ (C_I - C_{IV} , Fig. 1) and one connecting $\beta 3$ and the polypeptide segment connecting $\beta 4$ and $\beta 5$ (C_{II} - C_{III} , Fig. 1) are invariantly found in all CTLDs characterised so far. In the CTLD 3D-structure, these conserved secondary structure elements form a compact scaffold for a number of loops, which in the present context collectively are referred to as the "loop-region", protruding out from the core. These loops are in the primary structure of the CTLDs organised in two segments, loop segment A, LSA, and loop segment B, LSB. LSA represents the long polypeptide segment connecting $\beta 2$ and $\beta 3$ which often lacks regular secondary structure and contains up to four loops. LSB

represents the polypeptide segment connecting the β -strands $\beta 3$ and $\beta 4$. Residues in LSA, together with single residues in $\beta 4$, have been shown to specify the Ca^{2+} - and ligand-binding sites of several CTLDs, including that of tetranectin. E.g. mutagenesis studies, involving substitution of
5 single or a few residues, have shown, that changes in binding specificity, Ca^{2+} -sensitivity and/or affinity can be accommodated by CTLD domains [Weis and Drickamer (1996), Chiba et al. (1999), Graversen et al. (2000)].

10 As noted above, overall sequence similarities between CTLDs are often limited, as assessed e.g. by aligning a prospective CTLD sequence with the group of structure-characterized CTLDs presented in Fig. 1, using sequence alignment
15 procedures and analysis tools in common use in the field of protein science. In such an alignment, typically 22-30% of the residues of the prospective CTLD will be identical with the corresponding residue in at least one of the structure-characterized CTLDs. The sequence alignment shown in Fig. 1
20 was strictly elucidated from actual 3D structure data, so the fact that the polypeptide segments of corresponding structural elements of the framework also exhibit strong sequence similarities provide a set of direct sequence-structure signatures, which can readily be inferred from the sequence alignment.

25 The implication is that also CTLDs, for which precise 3D structural information is not yet available, can nonetheless be used as frameworks in the construction of new classes of CTLD libraries. The specific additional steps involved in preparing starting materials for the construction
30 of such a new class of CTLD library on the basis of a CTLD, for which no precise 3D structure is available, would be the following: (1) Alignment of the sequence of the new CTLD with the sequence shown in Fig. 1; and (2) Assignment of approximate locations of framework structural elements

as guided by the sequence alignment, observing any requirement for minor adjustment of the alignment to ensure precise alignment of the four canonical cysteine residues involved in the formation of the two conserved disulfide bridges (C_I-C_{IV} and $C_{II}-C_{III}$, in Fig. 1). The main objective of these steps would be to identify the sequence location of the loop-region of the new CTLD, as flanked in the sequence by segments corresponding to the $\beta 2$ -, $\beta 3$ - and $\beta 4$ -strands. To provide further guidance in this the results of an analysis of the sequences of 29 bona fide CTLDs are given in Table 1 below in the form of typical tetrapeptide sequences, and their consensus sequences, found as parts of CTLD $\beta 2$ - and $\beta 3$ -strands, and the precise location of the $\beta 4$ -strand by position and sequence characteristics as elucidated.

Table 1: $\beta 2$, $\beta 3$ and $\beta 4$ consensus elements analysis

CTLD	$\beta 2$	---	LSA	---	LSB	$\beta 3$	LSB	$\beta 4$
IX-A	W I G L R W - - - Q K V Q C N S E W S D G S S V S - - Y E N W I E - - - A E S K T - - - C L G L E K E T D F R K K V N I Y C							
MGL	W I G L T D Q - - - N G P - - - W H W V D G T D F E K G F K N W A P - - - L Q P D N W F G H G L G G G E D C A H I T T G - - G F W N D D V C							
LIT	W I G L H D P K K N R - - - W H W S S G S L V S - - Y K S W G I - - - G A P S N V P - - - G Y - C V S L T S S T G F Q K W K D V P C							
CHL	W I G L T D E N Q E G E - - - W Q W V D G T D T R S S F T F W K E - - - G E P N N R G F - - - N E D C A H V T S - - G Q W N D V Y C							
IGE-FCR	W I G L R N L D L K G E P I W V - - D G S H V D - - Y S N W A P - - - G E P T S R S Q - - - G E D C V M H R G S - - G R W N D A F C							
TCL-1	W I G L T D K D S E G T - - - W K W V D G T P P L T - - T A F W S T - - - D E P N D G A V N - - - G E D C V S L Y Y H T Q E F K N W N D L A C							
KUCR	W I G L T D Q G T E G N - - - W R W V D G T P P D V Q S R R F W R K - - - G Q P D W R H G N G E - - - R E D C V H L Q - - - R M W N D M A C							
CD94	W I G L S Y S E E H T A - - - W L W E N G S A L S Q - Y L S F E T - - - F N T K N - - - C I A Y N P N - - - G N A L D E S C							
CPCP	W I G L N D R T I E G D F R W S - - - D G H P M Q - - - F E N W R P - - - N Q P D N F F A A - - - G E D C V V M I W H E K G E W N D V P C							
PAP	W I G L H D P T O G T E P N G E G - - - W E W S S S D V M N - - - Y F A W E R - - - N - P S T I S S P G H - - - C A S L S R S T A F L R W K D V N C							
NEU	W I G L N D R I V E Q D - - - F Q W T D N T G L Q - - - Y E N W R E - - - N Q P D N F F A G - - - G E D C V V L V S H E I G K W N D V P C							
ESL	W I G I R K V N N V - - - W W W - V G T Q K P L T E A K N W A P - - - G E P N N R Q K - - - D E D C V E I Y I K R E K D V G H W N D E R C							
NK92A	W M G L S D L N Q E G T - - - W Q W V D G S P L L P S - - F K Q Y W N R - - - G E P N N V G - - - E E D C A E F S G N - - G W N D D K C							
GP120	W I G L F R N V - E G T - - - W L W I N N S P V S - - F V N W N T - - - G D P S G E - - - R N D C V A L H A S S - - G F W S N I B C							
MFR	W I G L N D M H A A E G T - - - W V D M T G A R I A Y K N W E T E I T - - - A Q P D G G K - - - T E N C A V L S G A A N G K W F E D K R C							
TN	W L G L S N V W N Q C N - - - W Q W S N A A M L R - - - Y K A W A E - - - E S Y - - - C V Y F K S T N - - N K W R S R A C							
SCGF	W L G V H D R R A E G L - - - Y L F E N G Q R V S - - F F A W H R S P R P E L G A Q S A S P H P L S P D Q P N G G T - - - L E N C V A Q A S D D - G S W N D D H C							
PLC	W L G A S D L N I E G R - - - W L W - E G Q R D Y E T G F K N W R P - - - E Q P D D W Y G H G L G G G E D C A H E T D D - - G R W N D D V C							
IX-B	W M G L B D - - - Q N G P - - - W K W V D G T D Y E T G F K N W R P - - - E S Y - - - C V Y F K S T N - - N K W R S R A C							
LY49A	W M G L S N V W N K K K D - - - W A W I D N R P S K L A L N T R K Y - - - N I R D G G - - - C M L I S K T - - - R L D N G N C							
TU14	W V G A D N - L Q D G A Y N F N W N D G V S L P T D S D L W S P - - - N E P S N P Q S W Q L - - - C V Q I W S K Y - N L L D D V G C							
ISP-A	Y L G M I E D Q T P G D - - - F H Y L D G A S V N - - - Y T N W Y P - - - G E P I G Q G - - - K E K C V E M Y T D - - G T W N D R G C							
BCON	Y L S M N D I S T E G R - - - F T Y P T G E I L V - - - Y S N W A D - - - G E P N N S D E G Q - - - P E N C L E I Y S D - - G K W N D V P C							
BCL43	Y L S M N D I S K E G K - - - F T Y P T G G S L D - - - Y S N W A P - - - G E P N N R A K D E G - - - P E N C L E I Y S D - - G N W N D I E C							
MBP-A	F L G I T D E V T E G Q - - - F M Y V T G G R L T - - - Y S N W K K - - - D E P N D H G S - - - G E D C V T I V D N - - G L W N D I S C							
SP-D	F L S M T D S K T E G K - - - F T Y P T G E S L V - - - Y S N W A P - - - G E P N D D G G - - - S E D C V E I F T N - - G K W N D R A C							
CL-L1	F I G V N D L E R E G Q - - - Y M F T D N T P L Q N - Y S N W N E - - - G E P S D P Y G - - - H E D C V E M L S S - - G R W N D T E C							
DCIR	F V G L S D P - - - E G Q R H W Q W V D Q T P - - - Y N E S S T E W H P - - - R E P S D P N - - - E R C V V L N F R K S P R W G - W N D V N C							

Notes: LSA, Loop Segment A; LSB, Loop Segment B.

Sequences taken from: Berglund and Petersen (1992) [TN, tetranectin]; Bertrand et al. (1996) [LIT, lithostatin]; Mann et al. (2000) [MGL, mouse macrophage galactose lectin, KUCR, Kupffer cell receptor, NEU, chicken neurocan, PLC, perlecan, HI-ASR, asialoglycoprotein receptor]; Mio et al. (1998) [CPCP, cartilage proteoglycan core protein, IGE-FCR, IgE Fc receptor, PAP, pancreatitis-associated protein, MFR, mouse macrophage receptor, NK92, Natural killer group, SCGF, stem cell growth factor]; Mizuno et al. (1997) [IX-A and B, factor IX/X binding protein, MBP, mannose binding protein]; Ohtani et al. (1999) [BCON, bovine conglutinin, BCL43, CL-L1, collectin liver 1, SP-A, surfactant protein A, SP-D, surfactant protein D]; Poget et al. (1999) [ESL, e-selectin, TU14, tunicate c-type lectin]; Tormo et al. (1999) [CD94, CD94 NK receptor domain, LY49A, LY49A NK receptor domain]; Zhang et al. (2000) [CHL, chicken hepatic lectin, TCL-1, trout c-type lectin, GP120, HIV gp 120-binding c-type lectin, DCIR, dendritic cell immuno-receptor]

Of the 29 β 2-strands,

14 were found to conform to the consensus sequence WIGX (of which 12 were WIGL sequences, 1 was a WIGI sequence and 1 was a WIGV sequence);

5 3 were found to conform to the consensus sequence WLGX (of which 1 was a WLGL sequence, 1 was a WLGV sequence and 1 was a WLGA sequence);

3 were found to be WMGL sequences;

10 3 were found to conform to the consensus sequence YLXM (of which 2 were YLSM sequences and 1 was an YLGM sequence);

2 were found to conform to the consensus sequence WVGX (of which 1 was a WVGL sequence and 1 was a WVGA sequence); and

15 the sequences of the remaining 4 β 2-strands in the collection were FLGI, FVGL, FIGV and FLSM sequences, respectively.

Therefore, it is concluded that the four-residue β 2 consensus sequence (" β 2cseq") may be specified as follows:

20 Residue 1: An aromatic residue, most preferably Trp, less preferably Phe and least preferably Tyr.

Residue 2: An aliphatic or non-polar residue, most preferably Ile, less preferably Leu or Met
25 and least preferably Val.

Residue 3: An aliphatic or hydrophilic residue, most preferably Gly and least preferably Ser.

Residue 4: An aliphatic or non-polar residue, most preferably Leu and less preferably Met, Val or Ile.

Accordingly the $\beta 2$ consensus sequence may be summarized as follows:

$\beta 2cseq: (\underline{W}, Y, F) - (\underline{I}, L, V, M) - (\underline{G}, S) - (\underline{L}, M, V, I),$

where the underlined residue denotes the most commonly found residue at that sequence position.

All 29 $\beta 3$ -strands analysed are initiated with the Cys₁₁ residue canonical for all known CTLD sequences, and of the 29 $\beta 3$ -strands,

5 were found to conform to the consensus sequence CVXI (of which 3 were CVEI sequences, 1 was a CVTI sequence and 1 was a CVQI sequence);

15 4 were found to conform to the consensus sequence CVXM (of which 2 were CVEM sequences, 1 was a CVVM sequence and 1 was a CVMM sequence);

6 were found to conform to the consensus sequence CVXL (of which 2 were CVVL sequences, 2 were a CVSL sequence, 1 was a CVHL sequence and 1 was CVAL sequence);

3 were found to conform to the consensus sequence CAXL (of which 2 were CAVL sequences and 1 was a CASL sequence);

25 2 were found to conform to the consensus sequence CAXF (of which 1 was 1 CAHF sequence and 1 was a CAEF sequence);

2 were found to conform to the consensus sequence CLXL (of which 1 was a CLEL sequence and 1 was a CLGL sequence); and

5 the sequences of the remaining 7 β 3-strands in the collection were CVYF, CVAQ, CAHV, CAHI, CLEI, CIAY, and CMLL sequences, respectively.

Therefore, it is concluded that the four-residue β 3 consensus sequence (" β 3cseq") may be specified as follows:

10 Residue 1: Cys, being the canonical Cys_{III} residue of CTLDs

Residue 2: An aliphatic or non-polar residue, most preferably Val, less preferably Ala or Leu and least preferably Ile or Met

15 Residue 3: Most commonly an aliphatic or charged residue, which most preferably is Glu

Residue 4: Most commonly an aliphatic, non-polar, or aromatic residue, most preferably Leu or Ile, less preferably Met or Phe and least preferably Tyr or Val.

20 Accordingly the β 3 consensus sequence may be summarized as follows:

β 3cseq: (C) - (V, A, L, I, M) - (E, X) - (L, I, M, F, Y, V),

where the underlined residue denotes the most commonly found residue at that sequence position.

25 It is observed from the known 3D-structures of CTLDs (Fig. 1), that the β 4-strands most often are comprised by five residues located in the primary structure at positions -6 to -2 relative to the canonical Cys_{III} residue of all known

CTLDs, and less often are comprised by four residues located at positions -5 to -2 relative to the canonical Cys_{III} residue of all known CTLDs. The residue located at position -3, relative to Cys_{III}, is involved in co-ordination of the site 2 calcium ion in CTLDs housing this site, and this notion is reflected in the observation, that of the 29 CTLD sequences analysed in Table 1, 27 have an Asp-residue or an Asn-residue at this position, whereas 2 CTLDs have a Ser at this position. From the known CTLD 3D-structures it is also noted, that the residue located at position -5, relative to the Cys_{III} residue, is involved in the formation of the hydrophobic core of the CTLD scaffold. This notion is reflected in the observation, that of the 29 CTLD sequences analysed 25 have a Trp-residue, 3 have a Leu-residue, and 1 an Ala-residue at this position. 18 of the 29 CTLD sequences analysed have an Asn-residue at position -4. Further, 19 of the 29 β 4-strand segments are preceded by a Gly residue.

Of the 29 central three residue motifs located at positions -5, -4 and -3 relative to the canonical Cys_{III} residue in the β 4-strand:

22 were of the sequence WXD (18 were WND, 2 were WKD, 1 was WFD and 1 was WWD),

2 were of the sequence WXN (1 was WVN and 1 was WSN),

and the remaining 5 motifs (WRS, LDD, LDN, LKS and ALD) were each represented once in the analysis.

It has now been found that each member of the family of CTLD domains represents an attractive opportunity for the construction of new protein libraries from which members with affinity for new ligand targets can be identified and isolated using screening or selection methods. Such libraries may be constructed by combining a CTLD framework struc-

ture in which the CTLD's loop-region is partially or completely replaced with one or more randomised polypeptide segments.

One such system, where the protein used as scaffold is tetranectin or the CTLD domain of tetranectin, is envisaged as a system of particular interest, not least because the stability of the trimeric complex of tetranectin protomers is very high (International Patent Application Publication No. WO 98/56906 A2).

- 10 Tetranectin is a trimeric glycoprotein [Holtet et al. (1997), Nielsen et al. (1997)], which has been isolated from human plasma and found to be present in the extracellular matrix in certain tissues. Tetranectin is known to bind calcium, complex polysaccharides, plasminogen, fi-
- 15 brinogen/fibrin, and apolipoprotein (a). The interaction with plasminogen and apolipoprotein (a) is mediated by the so-called kringle 4 protein domain therein. This interaction is known to be sensitive to calcium and to derivatives of the amino acid lysine [Graversen et al. (1998)].
- 20 A human tetranectin gene has been characterised, and both human and murine tetranectin cDNA clones have been isolated. Both the human and the murine mature protein comprise 181 amino acid residues (Fig. 2). The 3D-structures of full length recombinant human tetranectin and of the
- 25 isolated tetranectin CTLD have been determined independently in two separate studies [Nielsen et al. (1997) and Kastrup et al. (1998)]. Tetranectin is a two- or possibly three-domain protein, i.e. the main part of the polypeptide chain comprises the CTLD (amino acid residues Gly53 to
- 30 Val181), whereas the region Leu26 to Lys52 encodes an alpha-helix governing trimerisation of the protein via the formation of a homotrimeric parallel coiled coil. The polypeptide segment Glu1 to Glu25 contains the binding site for

complex polysaccharides (Lys6 to Lys15) [Lorentsen et al. (2000)] and appears to contribute to stabilisation of the trimeric structure [Holtet et al. (1997)]. The two amino acid residues Lys148 and Glu150, localised in loop 4, and Asp165 (localised in β 4) have been shown to be of critical importance for plasminogen kringle 4 binding, whereas the residues Ile140 (in loop 3) and Lys166 and Arg167 (in β 4) have been shown to be of some importance [Graversen et al. (1998)]. Substitution of Thr149 (in loop 4) with an aromatic residue has been shown to significantly increase affinity of tetranectin to kringle 4 and to increase affinity for plasminogen kringle 2 to a level comparable to the affinity of wild type tetranectin for kringle 4 [Graversen et al. (2000)].

15 OBJECT OF THE INVENTION

The object of the invention is to provide a new practicable method for the generation of useful protein products endowed with binding sites able to bind substance of interest with high affinity and specificity.

20 The invention describes one way in which such new and useful protein products may advantageously be obtained by applying standard combinatorial protein chemistry methods, commonly used in the recombinant antibody field, to generate randomised combinatorial libraries of protein modules, 25 in which each member contains an essentially common core structure similar to that of a CTLD.

The variation of binding site configuration among naturally occurring CTLDs shows that their common core structure can accommodate many essentially different configurations of the ligand binding site. CTLDs are therefore particularly well suited to serve as a basis for constructing such new and useful protein products with desired binding properties.

In terms of practical application, the new artificial CTLD protein products can be employed in applications in which antibody products are presently used as key reagents in technical biochemical assay systems or medical *in vitro* or
5 *in vivo* diagnostic assay systems or as active components in therapeutic compositions.

In terms of use as components of *in vitro* assay systems, the artificial CTLD protein products are preferable to antibody derivatives as each binding site in the new protein
10 product is harboured in a single structurally autonomous protein domain. CTLD domains are resistant to proteolysis, and neither stability nor access to the ligand-binding site is compromised by the attachment of other protein domains to the N- or C-terminus of the CTLD. Accordingly, the CTLD
15 binding module may readily be utilized as a building block for the construction of modular molecular assemblies, e.g. harbouring multiple CLTDs of identical or nonidentical specificity in addition to appropriate reporter modules like peroxidases, phosphatases or any other signal-
20 mediating moiety.

In terms of *in vivo* use as essential component of compositions to be used for *in vivo* diagnostic or therapeutic purposes, artificial CTLD protein products constructed on the basis of human CTLDs are virtually identical to the corresponding natural CTLD protein already present in the body,
25 and are therefore expected to elicit minimal immunological response in the patient. Single CTLDs are about half the mass of the smallest functional antibody derivative, the single-chain Fv derivative, and this small size may in some
30 applications be advantageous as it may provide better tissue penetration and distribution, as well as a shorter half-life in circulation. Multivalent formats of CTLD proteins, e.g. corresponding to the complete tetranectin trimer or the further multimerized collectins, like e.g.

mannose binding protein, provide increased binding capacity and avidity and longer circulation half-life.

One particular advantage of the preferred embodiment of the invention, arises from the fact that mammalian tetranectins, as exemplified by murine and human tetranectin, are of essentially identical structure. This conservation among species is of great practical importance as it allows straightforward swapping of polypeptide segments defining ligand-binding specificity between e.g. murine and human tetranectin derivatives. The option of facile swapping of species genetic background between tetranectin derivatives is in marked contrast to the well-known complications of effecting the "humanisation" of murine antibody derivatives.

Further advantages of the invention are:

The availability of a general and simple procedure for reliable conversion of an initially selected protein derivative into a final protein product, which without further reformatting may be produced in bacteria (e.g. *Escherichia coli*) both in small and in large scale (International Patent Application Publication No. WO 94/18227 A2).

The option of including several identical or non-identical binding sites in the same functional protein unit by simple and general means, thereby enabling the exploitation even of weak affinities by means of avidity in the interaction, or the construction of bi- or heterofunctional molecular assemblies (International Patent Application Publication No. WO 98/56906 A2).

The possibility of modulating binding by addition or removal of divalent metal ions (e.g. calcium ions) in combination with one or more preserved metal binding site(s) in the CTLDs.

SUMMARY OF THE INVENTION

The present invention provides a great number of novel and useful proteins each being a protein having the scaffold structure of C-type lectin-like domains (CTLD), said protein comprising a variant of a model CTLD wherein the α -helices and β -strands and connecting segments are conserved to such a degree that the scaffold structure of the CTLD is substantially maintained, while the loop region is altered by amino acid substitution, deletion, insertion or any combination thereof, with the proviso that said protein is not any of the known CTLD loop derivatives of C-type lectin-like proteins or C-type lectins listed in the following Table 2.

TABLE 2: Known $\beta 2$, $\beta 3$, $\beta 4$, LSA and LSB CTLD derivatives
 Table 2A: LSA derivatives ($\beta 2$ and $\beta 3$ consensus elements are underlined)

CTLD	Mut.	LSA sequence (one letter code)	Reference
HTN	TND116A	<u>W L G L</u> N A M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T E N C A V L	Graversen et al. (1998)
	TNE120A	<u>W L G L</u> N D M A A A G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T E N C A V L	Graversen et al. (1998)
	TNK134A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y A N W E T E I T A Q P D G G K T E N C A V L	Graversen et al. (1998)
	TNT140A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E A T A Q P D G G K T E N C A V L	Graversen et al. (1998)
	TNQ143A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A A P D G G K T E N C A V L	Graversen et al. (1998)
	TND145A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P A G G K T E N C A V L	Graversen et al. (1998)
	TNK148A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G A T E N C A V L	Graversen et al. (1998)
	TNK148M	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G M T E N C A V L	Graversen et al. (2000)
	TNK148R	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G R T E N C A V L	Graversen et al. (2000)
	TNT149F	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K F E N C A V L	Graversen et al. (2000)
	TNT149M	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K M E N C A V L	Graversen et al. (2000)
	TNT149R	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K R E N C A V L	Graversen et al. (2000)
	TNT149Y	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K Y E N C A V L	Graversen et al. (2000)
	TNE150A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T A N C A V L	Graversen et al. (1998)
	TNE150D	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T D N C A V L	Graversen et al. (2000)
	TNE150Q	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T E A C A V L	Graversen et al. (1998)
	TNN151A	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K T Q N C A V L	Graversen et al. (2000)
	TNK148R, T149Y	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K Y E N C A V L	Graversen et al. (1998)
	TNT149Y, E150Q	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K Y E N C A V L	Graversen et al. (2000)
	TNT149Y, D165N	<u>W L G L</u> N D M A A E G T W V D M T G A R I A Y K N W E T E I T A Q P D G G K Y E N C A V L	Graversen et al. (2000)
rMBP	QPD	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D H G S G E D C V T I	Drickamer (1992)
	N187D	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P D D H G S G E D C V T I	Iobst et al. (1994)
	H189A	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P N D A G S G E D C V T I	Iobst et al. (1994)
	H189G	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P N D G S G E D C V T I	Iobst et al. (1994)
	QPDW	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W G S G E D C V T I	Iobst & Drickamer (1994)
	QPDWG	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G H L G G G E D C V T I	Iobst & Drickamer (1994)
	QPDWG/Y/A	<u>F L G I</u> T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W A G H L G G G E D C V T I	Iobst & Drickamer (1994)

QPDWG/Y/Q	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Q G H L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/G/A	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y A H L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/H/A	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G A L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/H/Q	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G Q L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/H/E	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G E L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/H/Y	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G Y L G G G E D C V T I	Iobst & Drickamer (1994)
QPDWG/-/G	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G H L G G G E D C V T I	Iobst & Drickamer (1994)
QPDF	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D F G S G E D C V T I	Iobst & Drickamer (1994)
QPDFG	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D F Y G H L G G G E D C V T I	Iobst & Drickamer (1994)
REGION 1	FLG I R K V N N V F M Y V T G G R L T Y S N W K K D E P N D H G S G E D C V T I	Blanck et al. (1996)
REGION 2	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P N R Q K D E D C V T I	Blanck et al. (1996)
RES. 189	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P N D G S G E D C V T I	Torgersen et al. (1998)
RES. 197	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D E P N D H G S G E D C V E I	Torgersen et al. (1998)
LOOP 3E	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W A P E Z P N D H G S G E D C V T I	Torgersen et al. (1998)
LOOP 3P	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W A D N E P N D H G S G E D C V T I	Torgersen et al. (1998)
REGION 4	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G H L G G G E D C V H I	Kolatkar et al. (1998)
REGION 4'	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W R P Q P D D W Y G H L G G G E D C V H I	Kolatkar et al. (1998)
QPDWG/QNG	FLG I T D Q N G Q F M Y V T G G R L T Y S N W K K D Q P D D W Y G H L G G G E D C V T I	Wragg & Drickamer (1999)
QPDWG/QNGP	FLG I T D Q N G P F M Y V T G G R L T Y S N W K K D Q P D D W Y G H L G G G E D C V T I	Wragg & Drickamer (1999)
MBP/CHL189	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K E Z E P N R G S G E D C V T I	Burrows et al. (1997)
MBP/CHL192	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K E Z E P N R G F N E D C V T I	Burrows et al. (1997)
MBP/CHL208	FLG I T D E V T E G Q F M Y V T G G R L T Y S N W K E Z E P N R G F N E D C A H V	Burrows et al. (1997)
rSP-A E195Q, R197D	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G Q P D G Q G K E K C V E M	McCormack et al. (1994)
AM2	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P R G Q G K E K C V T I	Honma et al. (1997)
AM3	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P N D H G S G E D C V T I	Honma et al. (1997)
E195A	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G A P R G Q G K E K C V E M	McCormack et al. (1997)
R197G	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P G G Q G K E K C V E M	McCormack et al. (1997)
E202A	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P R G Q G K A K C V E M	McCormack et al. (1997)
N187S	Y L G M I E D Q T P G D F H Y L D G A S V S Y T N W Y P G E P R G Q G K E K C V E M	McCormack et al. (1997)
R197A	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P A G Q G K E K C V E M	Pattanaajitvilai et al. (1998)
R197K	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P K G Q G K E K C V E M	Pattanaajitvilai et al. (1998)
R197H	Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P H G Q G K E K C V E M	Pattanaajitvilai et al. (1998)

R197D	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P D G Q G K E K C V E M</u>	Pattana jittvilai et al. (1998)
R197N	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P N G Q G K E K C V E M</u>	Pattana jittvilai et al. (1998)
E195Q	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G Q P R G Q G K E K C V E M</u>	Tsunezawa et al. (1998)
K201A	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P R G Q G A E K C V E M</u>	Tsunezawa et al. (1998)
K203A	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P R G Q G K E A C V E M</u>	Tsunezawa et al. (1998)
E197A, K201A, K203A	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G A P R G Q G A E A C V E M</u>	Tsunezawa et al. (1998)
ad3	<u>Y L G M I E D Q T P G D F H Y L D G A S V N Y T N W Y P G E P N N N G G A E N C V E I</u>	Sano et al. (1998)
ad4	<u>Y L G M I E D Q T E G K F T Y P T G E A L V Y S N W A P G E P N N G G A E N C V E I</u>	Sano et al. (1998)
rat ama4	<u>Y L G M I E D Q T E G Q F M Y V T G G R L T Y S N W K K D E P R G Q G K E K C V E M</u>	Chiba et al. (1999)
hsp-A R199A	<u>Y V G L T E G P S P G D F R Y S D G T P V N Y T N W Y R G E P A G A G K E Q C V E M</u>	Tsunezawa et al. (1998)
K201A	<u>Y V G L T E G P S P G D F R Y S D G T P V N Y T N W Y R G E P A G R G A E Q C V E M</u>	Tsunezawa et al. (1998)
hum ama4	<u>Y V G L T E G P T E G Q F M Y V T G G R L T Y S N W K K D E P R G R G K E Q C V E M</u>	Chiba et al. (1999)
rSP-D E321Q, N323D	<u>F L S M T D V G T E G K F T Y P T G E A L V Y S N W A P G Q P D N N G G A E N C V E I</u>	Ogasawara & Voelker (1995)
h-esl K67A	<u>W I G I R K V N N V W V V G T Q A P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Erbe et al.
K74A	<u>W I G I R K V N N V W V V G T Q K P L T E E A A N W A P G E P N N R Q K D E D C V E I</u>	Erbe et al.
R84A, K86A	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N A Q A D E D C V E I</u>	Erbe et al.
R84A	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N A Q K D E D C V E I</u>	Kogan et al. (1995)
R84K	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N A Q K D E D C V E I</u>	Kogan et al. (1995)
R84K, D89G	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N A Q K D E D C V E I</u>	Kogan et al. (1995)
A77K	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
A77K, P78K	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
A77K, P78K, R84A	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N A Q K D E D C V E I</u>	Kogan et al. (1995)
D87E	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
D87N	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
D89N	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
D89E	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
A77K, E80Q, N82D	<u>W I G I R K V N N V W V V G T Q K P L T E E A K N W A P G E P N N R Q K D E D C V E I</u>	Kogan et al. (1995)
A77K	<u>W I G I R K N N K T W T W V G T K K A L T N E A E N W K D N E P N N K R N N E D C V E I</u>	Revelle et al. (1996)
A77K, E80D, N82D	<u>W I G I R K N N K T W T W V G T K K A L T N E A E N W K D N Q P D N K R N N E D C V E I</u>	Revelle et al. (1996)
2A/R	<u>W I G L T D Q N G P W R W V D G T D Y E K G F T H W R P K Q P D N W Y G H G L G G E D C A H F</u>	Iobst & Drickamer (1996)
2K/G	<u>W I G L T D Q N G P W R W V D G T D Y E K G F T H W A P G Q P D N W Y G H G L G G E D C A H F</u>	Iobst & Drickamer (1996)
2A/R, 2K/G	<u>W I G L T D Q N G P W R W V D G T D Y E K G F T H W R P G Q P D N W Y G H G L G G E D C A H F</u>	Iobst & Drickamer (1996)

4F/I	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A H I	Iobst & Drickamer (1996)
4H/A	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A A F	Iobst & Drickamer (1996)
4H/E	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A E F	Iobst & Drickamer (1996)
4H/Q	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A Q F	Iobst & Drickamer (1996)
4H/N	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A N F	Iobst & Drickamer (1996)
4H/Y	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A Y F	Iobst & Drickamer (1996)
4H/D	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A D F	Iobst & Drickamer (1996)
4H/K	WIGL T D Q N G P W R W V D G T D Y E K G F T H W A P K Q P D N W Y G H G L G G G E D C A K F	Iobst & Drickamer (1996)
2A/R, 2K/G, 4H/A	WIGL T D Q N G P W R W V D G T D Y E K G F T H W R P G Q P D N W Y G H G L G G G E D C A A F	Iobst & Drickamer (1996)
RBL	WIGL T D Q N G P W K W V D G T D Y E T G F K N W R P G Q P D D W Y G H G L G G G E D C A A F	Iobst & Drickamer (1996)
CHL	W I G L T D E N Q E G E W Q W V D G T D T R S S F T F W K E G E P N N A G F N E D C A H V	Burrows et al. (1997)
	W I G L T D E N Q E G E W Q W V D G T D T R S S F T F W K E G E P N N R A F N E D C A H V	Burrows et al. (1997)
	W I G L T D E N Q E G E W Q W V D G T D T R S S F T F W K E G E P N N R G A N E D C A H V	Burrows et al. (1997)
	W I G L T D E N Q E G E W Q W V D G T D T R S S F T F W K E G E P N N R G F A E D C A H V	Burrows et al. (1997)

Table 2B: LSB derivatives ($\beta 3$ and $\beta 4$ consensus elements are underlined)

CTL	Mut.	LSB sequence (one letter code)	Reference
HTN	TNK163A	<u>C A V L S G A A N G A W F D K R C</u>	Graversen et al. (1998)
	TNK166A	<u>C A V L S G A A N G K W F D A R C</u>	Graversen et al. (1998)
	TNR167A	<u>C A V L S G A A N G K W F D K A C</u>	Graversen et al. (1998)
	TNF164L	<u>C A V L S G A A N G K W L D K R C</u>	Graversen et al. (1998)
	TND165A	<u>C A V L S G A A N G K W F A K R C</u>	Graversen et al. (1998)
	TND165E	<u>C A V L S G A A N G K W F E K R C</u>	Graversen et al. (2000)
	TND165N	<u>C A V L S G A A N G K W F N K R C</u>	Graversen et al. (2000)
rMBP	I207V	<u>C V T I V D N G L W N D V S C</u>	Iobst et al. (1994)
	I207L	<u>C V T I V D N G L W N D L S C</u>	Iobst et al. (1994)
	I207A	<u>C V T I V D N G L W N D A S C</u>	Iobst et al. (1994)
	I207E	<u>C V T I V D N G L W N D E S C</u>	Torgensen et al. (1996)
Region 4E		<u>C V T I V Y I K R E K D N G L W N D I S C</u>	Torgensen et al. (1996)
Region 4P		<u>C V T I V Y I K S P S D N G L W N D I S C</u>	Torgensen et al. (1996)
207VY		<u>C V T I V D N G L W N D V Y C</u>	Burrows et al. (1997)
$\beta 34$		<u>C A H V W T S G Q W N D V Y C</u>	Burrows et al. (1997)
h-es1	Y94F	<u>C V E I F I K R E K D V G M W N D E R C</u>	Kogan et al. (1995)
	Y94R	<u>C V E I R I K R E K D V G M W N D E R C</u>	Kogan et al. (1995)
	Y94D	<u>C V E I D I K R E K D V G M W N D E R C</u>	Kogan et al. (1995)
	Y94A	<u>C V E I A I K R E K D V G M W N D E R C</u>	Kogan et al. (1995)
	Y94S	<u>C V E I S I K R E K D V G M W N D E R C</u>	Kogan et al. (1995)
	E107D	<u>C V E I Y I K R E K D V G M W N D D R C</u>	Kogan et al. (1995)
	E107A	<u>C V E I Y I K R E K D V G M W N D A R C</u>	Kogan et al. (1995)
	E107N	<u>C V E I Y I K R E K D V G M W N D N R C</u>	Kogan et al. (1995)
	E107K	<u>C V E I Y I K R E K D V G M W N D K R C</u>	Kogan et al. (1995)
	E107Q	<u>C V E I Y I K R E K D V G M W N D Q R C</u>	Kogan et al. (1995)
	R97D	<u>C V E I Y I K D E K D V G M W N D E R C</u>	Revelle et al. (1996)
	R97S	<u>C V E I Y I K S E K D V G M W N D E R C</u>	Revelle et al. (1996)

R97E	<u>C V E I Y I K E E K D V G M W N D E R C</u>	Revelle et al. (1996)
h-ps1 K96Q	<u>C V E I Y I Q S P S A P G M W N D E H C</u>	Revelle et al. (1996)
K96R	<u>C V E I Y I R S P S A P G M W N D E H C</u>	Revelle et al. (1996)
K96E	<u>C V E I Y I E S P S A P G M W N D E H C</u>	Revelle et al. (1996)
S97A	<u>C V E I Y I K A P S A P G M W N D E H C</u>	Revelle et al. (1996)
S97D	<u>C V E I Y I K D P S A P G M W N D E H C</u>	Revelle et al. (1996)
S97R	<u>C V E I Y I K R P S A P G M W N D E H C</u>	Revelle et al. (1996)
REK	<u>C V E I Y I K R E K A P G M W N D E H C</u>	Revelle et al. (1996)
S99D	<u>C V E I Y I K S P D A P G M W N D E H C</u>	Revelle et al. (1996)
CHL V191A	<u>C A H V W T S G Q W N D A Y C</u>	Burrows et al. (1997)
Y192A	<u>C A H V W T S G Q W N D V A C</u>	Burrows et al. (1997)

2C: Other TN CTLD derivatives

CTLD	Mut.	TN sequence (one letter code)	Reference
hTN	TNR169A	S G A A N G K W F D K R C A D Q	Graversen et al. (1998)
	TNS85C	C I S R G G T L G T P Q T	Jaquinod et al. (1999)
Notes:			
hTN: human tetranectin;			
rMBP: rat mannanose binding protein,			
rSP-A: rat surfactant protein-A,			
hSP-A: human surfactant protein-A,			
rSP-D: rat surfactant protein-D;			
h-es1: human e-selectin;			
h-ps1: human p-selectin;			
MGR: macrophage galactose receptor;			
REL: rat hepatic lectin,			
CHL: chicken hepatic lectin			

Normally the model CTLD is defined by having a 3D structure that conforms to the secondary-structure arrangement illustrated in Fig. 1 characterized by the following main secondary structure elements:

- 5 five β -strands and two α -helices sequentially appearing in the order $\beta 1$, $\alpha 1$, $\alpha 2$, $\beta 2$, $\beta 3$, $\beta 4$, and $\beta 5$, the β -strands being arranged in two anti-parallel β -sheets, one composed of $\beta 1$ and $\beta 5$, the other composed of $\beta 2$, $\beta 3$ and $\beta 4$,
 - 10 at least two disulfide bridges, one connecting $\alpha 1$ and $\beta 5$ and one connecting $\beta 3$ and the polypeptide segment connecting $\beta 4$ and $\beta 5$,
 - 15 a loop region consisting of two polypeptide segments, loop segment A (LSA) connecting $\beta 2$ and $\beta 3$ and comprising typically 15-70 or, less typically, 5-14 amino acid residues, and loop segment B (LSB) connecting $\beta 3$ and $\beta 4$ and comprising typically 5-12 or less typically, 2-4 amino acid residues.
- However, also a CTLD, for which no precise 3D structure
- 20 is available, can be used as a model CTLD, such CTLD being defined by showing sequence similarity to a previously recognised member of the CTLD family as expressed by an amino acid sequence identity of at least 22 %, preferably at least 25 % and more preferably at least 30
 - 25 %, and by containing the cysteine residues necessary for establishing the conserved two-disulfide bridge topology (i.e. Cys_I, Cys_{II}, Cys_{III} and Cys_{IV}). The loop region, consisting of the loop segments LSA and LSB, and its flanking β -strand structural elements can then be identified
 - 30 by inspection of the sequence alignment with the collection of CTLDs shown in Fig. 1, which provides identification of the sequence locations of the $\beta 2$ - and $\beta 3$ -strands

with the further corroboration provided by comparison of these sequences with the four-residue consensus sequences, $\beta 2cseq$ and $\beta 3cseq$, and the $\beta 4$ strand segment located typically at positions -6 to -2 and less typically at positions -5 to -2 relative to the conserved Cys_{III} residue and with the characteristic residues at positions -5 and -3 as elucidated from Table 1 and deducted above under BACKGROUND OF THE INVENTION.

The same considerations apply for determining whether in a model CTLD the α -helices and β -strands and connecting segments are conserved to such a degree that the scaffold structure of the CTLD is substantially maintained.

It may be desirable that up to 10, preferably up to 4, and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and/or β -strands and/or connecting segments of the model CTLD. In particular, changes of up to 4 residues may be made in the β -strands of the model CTLD as a consequence of the introduction of recognition sites for one or more restriction endonucleases in the nucleotide sequence encoding the CTLD to facilitate the excision of part or all of the loop region and the insertion of an altered amino acid sequence instead while the scaffold structure of the CTLD is substantially maintained.

Of particular interest are proteins wherein the model CTLD is that of a tetranectin. Well known tetranectins the CTLDs of which can be used as model CTLDs are human tetranectin and murine tetranectin. The proteins according to the invention thus comprise variants of such model CTLDs.

The proteins according to the invention may comprise N-terminal and/or C-terminal extensions of the CTLD vari-

ant, and such extensions may for example contain effector, enzyme, further binding and/or multimerising functions. In particular, said extension may be the non-CTLD-
portions of a native C-type lectin-like protein or C-type
5 lectin or a "soluble" variant thereof lacking a functional transmembrane domain.

The proteins according to the invention may also be multimers of a moiety comprising the CTLD variant, e.g. derivatives of the native tetranectin trimer.

10 In a preferred aspect the present invention provides a combinatorial library of proteins having the scaffold structure of C-type lectin-like domains (CTLD), said proteins comprising variants of a model CTLD wherein the α -helices and β -strands are conserved to such a degree that
15 the scaffold structure of the CTLD is substantially maintained, while the loop region or parts of the loop region of the CTLD is randomised with respect to amino acid sequence and/or number of amino acid residues.

The proteins making up such a library comprise variants
20 of model CTLDs defined as for the above proteins according to the invention, and the variants may include the changes stated for those proteins.

In particular, the combinatorial library according to the invention may consist of proteins wherein the model CTLD
25 is that of a tetranectin, e.g. that of human tetranectin or that of murine tetranectin.

The combinatorial library according to the invention may consist of proteins comprising N-terminal and/or C-terminal extensions of the CTLD variant, and such extensions may for example contain effector, enzyme, further
30 binding and/or multimerising functions. In particular,

said extensions may be the non-CTLD-portions of a native C-type lectin-like protein or C-type lectin or a "soluble" variant thereof lacking a functional transmembrane domain.

- 5 The combinatorial library according to the invention may also consist of proteins that are multimers of a moiety comprising the CTLD variant, e.g. derivatives of the native tetranectin trimer.

The present invention also provides derivatives of a native tetranectin wherein up to 10, preferably up to 4,
10 and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and/or β -strands and/or connecting segments of its CTLD as well as nucleic acids encoding such derivatives. Specific derivatives appear from SEQ ID Nos: 02, 04, 09, 11, 13, 15, 29,
15 31, 36, and 38; and nucleic acids comprising nucleotide inserts encoding specific tetranectin derivatives appear from SEQ ID Nos: 12, 14, 35, and 37.

The invention comprises a method of constructing a tetranectin derivative adapted for the preparation of a combinatorial library according to the invention, wherein the nucleic acid encoding the tetranectin derivative has been modified to generate endonuclease restriction sites within nucleic acid segments encoding β 2, β 3 or β 4, or
20 up to 30 nucleotides upstream or downstream in the sequence from any nucleotide which belongs to a nucleic acid segment encoding β 2, β 3 or β 4.
25

The invention also comprises the use of a nucleotide sequence encoding a tetranectin, or a derivative thereof
30 wherein the scaffold structure of its CTLD is substantially maintained, for preparing a library of nucleotide sequences encoding related proteins by randomising part

or all of the nucleic acid sequence encoding the loop region of its CTLD.

Further, the present invention provides nucleic acid comprising any nucleotide sequence encoding a protein according to the invention.

In particular, the invention provides a library of nucleic acids encoding proteins of a combinatorial library according to the invention, in which the members of the ensemble of nucleic acids, that collectively constitute said library of nucleic acids, are able to be expressed in a display system, which provides for a logical, physical or chemical link between entities displaying phenotypes representing properties of the displayed expression products and their corresponding genotypes.

In such a library the display system may be selected from

(I) a phage display system such as

(1) a filamentous phage fd in which the library of nucleic acids is inserted into

(a) a phagemid vector,

(b) the viral genome of a phage

(c) purified viral nucleic acid in purified single- or double-stranded form, or

(2) a phage lambda in which the library is inserted into

(a) purified phage lambda DNA, or

(b) the nucleic acid in lambda phage particles; or

(II) a viral display system in which the library of nucleic acids is inserted into the viral nucleic acid of a eukaryotic virus such as baculovirus; or

- (III) a cell-based display system in which the library of nucleic acids is inserted into, or adjoined to, a nucleic acid carrier able to integrate either into the host genome or into an extrachromosomal element able to maintain and express itself within the cell and suitable for cell-surface display on the surface of
- (a) bacterial cells,
 - (b) yeast cells, or
 - (c) mammalian cells; or
- (IV) a nucleic acid entity suitable for ribosome linked display into which the library of nucleic acid is inserted; or
- (V) a plasmid suitable for plasmid linked display into which the library of nucleic acid is inserted.

A well-known and useful display system is the "Recombinant Phage Antibody System" with the phagemid vector "pCANTAB 5E" supplied by Amersham Pharmacia Biotech (code no. 27-9401-01).

Further, the present invention provides a method of preparing a protein according to the invention, wherein the protein comprises at least one or more, identical or not identical, CTLD domains with novel loop-region sequences which has (have) been isolated from one or more CTLD libraries by screening or selection. At least one such CTLD domain may have been further modified by mutagenesis; and the protein containing at least one CTLD domain may have been assembled from two or more components by chemical or enzymatic coupling or crosslinking.

Also, the present invention provides a method of preparing a combinatorial library according to the invention comprising the following steps:

- 1) inserting nucleic acid encoding a protein comprising a model CTLD into a suitable vector,
- 2) if necessary, introducing restriction endonuclease recognition sites by site directed mutagenesis,
5 said recognition sites being properly located in the sequence at or close to the ends of the sequence encoding the loop region of the CTLD or part thereof,
- 3) excising the DNA fragment encoding the loop region
10 or part thereof by use of the proper restriction endonucleases,
- 4) ligating mixtures of DNA fragments into the restricted vector, and
- 5) inducing the vector to express randomised proteins
15 having the scaffold structure of CTLDs in a suitable medium.

In a further aspect, the present invention provides a method of screening a combinatorial library according to the invention for binding to a specific target which comprises the following steps:

20

- 1) expressing a nucleic acids library according to any one of claims 59-61 to display the library of proteins in the display system;
- 2) contacting the collection of entities displayed
25 with a suitably tagged target substance for which isolation of a CTLD-derived exhibiting affinity for said target substance is desired;
- 3) harvesting subpopulations of the entities displayed that exhibit affinity for said target substance by
30 means of affinity-based selective extractions, utilizing the tag to which said target substance is conjugated or physically attached or adhering to as a vehicle or means of affinity purification, a procedure commonly referred to in the field as "affin-

- ity panning", followed by re-amplification of the sub-library;
- 4) isolating progressively better binders by repeated rounds of panning and re-amplification until a suitably small number of good candidate binders is obtained; and,
- 5) if desired, isolating each of the good candidates as an individual clone and subjecting it to ordinary functional and structural characterisation in preparation for final selection of one or more preferred product clones.

In a still further aspect, the present invention provides a method of reformatting a protein according to the invention or selected from a combinatorial library according to the invention and containing a CTLD variant exhibiting desired binding properties, in a desired alternative species-compatible framework by excising the nucleic acid fragment encoding the loop region-substituting polypeptide and any required single framework mutations from the nucleic acid encoding said protein using PCR technology, site directed mutagenesis or restriction enzyme digestion and inserting said nucleic acid fragment into the appropriate location(s) in a display- or protein expression vector that harbours a nucleic acid sequence encoding the desired alternative CTLD framework.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows an alignment of the amino acid sequences of ten CTLDs of known 3D-structure. The sequence locations of main secondary structure elements are indicated above each sequence, labelled in sequential numerical order as " α N", denoting α -helix number N, and " β M", denoting β -strand number M.

The four cysteine residues involved in the formation of the two conserved disulfide bridges of CTLDs are indicated and enumerated in the Figure as "C_I", "C_{II}", "C_{III}" and "C_{IV}", respectively. The two conserved disulfide
5 bridges are C_I-C_{IV} and C_{II}-C_{III}, respectively.

The ten C-type lectins are

- hTN: human tetranectin [Nielsen et al. (1997)];
- MBP: mannose binding protein [Weis et al. (1991); Sheriff et al. (1994)];
- 10 SP-D: surfactant protein D [Håkansson et al. (1999)];
- LY49A: NK receptor LY49A [Tormo et al. (1999)];
- H1-ASR: H1 subunit of the asialoglycoprotein receptor [Meier et al. (2000)];
- MMR-4: macrophage mannose receptor domain 4 [Feinberg et
15 al. (2000)];
- IX-A and IX-B: coagulation factors IX/X-binding protein domain A and B, respectively [Mizuno et al. (1997)];
- Lit: lithostatine [Bertrand et al. (1996)];
- 20 TU14: tunicate C-type lectin [Poget et al. (1999)].

Fig. 2 shows an alignment of the nucleotide and amino acid sequences of the coding regions of the mature forms of human and murine tetranectin with an indication of known secondary structural elements.

- 25 hTN: human tetranectin; nucleotide sequence from Berglund and Petersen (1992).

mTN: murine tetranectin; nucleotide sequence from Sørensen et al. (1995).

Secondary structure elements from Nielsen et al. (1997).

" α " denotes an α -helix; " β " denotes a β -strand; and "L"

5 denotes a loop.

Fig. 3 shows an alignment of the nucleotide and amino acid sequences of human and murine tlec coding regions.

htlec: the sequence derived from hTN; mtlec: the sequence derived from mTN. The position of the restriction endonuclease sites for *Bgl* II, *Kpn* I, and *Mun* I are indicated.

Fig. 4 shows an alignment of the nucleotide and amino acid sequences of human and murine tCTLD coding regions.

htCTLD: the sequence derived from hTN; mtCTLD: the sequence derived from mTN. The position of the restriction endonuclease sites for *Bgl* II, *Kpn* I, and *Mun* I are indicated.

Fig. 5 shows an outline of the pT7H6FX-htlec expression plasmid. The FX-htlec fragment was inserted into pT7H6 [Christensen et al. (1991)] between the Bam HI and Hind III cloning sites.

Fig. 6 shows the amino acid sequence (one letter code) of the FX-htlec part of the H6FX-htlec fusion protein produced by pT7H6FX-htlec.

Fig. 7 shows an outline of the pT7H6FX-htCTLD expression plasmid. The FX-htCTLD fragment was inserted into pT7H6 [Christensen et al. (1991)] between the Bam HI and Hind III cloning sites.

Fig. 8 shows the amino acid sequence (one letter code) of the FX-htCTLD part of the H6FX-htCTLD fusion protein produced by pT7H6FX-htCTLD.

Fig. 9 shows an outline of the pPhTN phagemid. The PhTN fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.

- 5 Fig. 10 shows the amino acid sequence (one letter code) of the PhTN part of the PhTN-gene III fusion protein produced by pPhTN.

- Fig. 11 shows an outline of the pPhTN3 phagemid. The PhTN3 fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.
- 10

Fig. 12 shows the amino acid sequence (one letter code) of the PhTN3 part of the PhTN3-gene III fusion protein produced by pPhTN3.

- 15 Fig. 13 shows an outline of the pPhTlec phagemid. The PhTlec fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.

- Fig. 14 shows the amino acid sequence (one letter code) of the PhTlec part of the PhTlec-gene III fusion protein produced by pPhTlec.
- 20

- Fig. 15 shows an outline of the pPhtCTLD phagemid. The PhtCTLD fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.
- 25

Fig. 16 shows the amino acid sequence (one letter code) of the PhtCTLD part of the PhtCTLD-gene III fusion protein produced by pPhtCTLD.

Fig. 17 shows an outline of the pUC-mtlec.

Fig. 18 shows an outline of the pT7H6FX-mtlec expression plasmid. The FX-mtlec fragment was inserted into pT7H6 [Christensen et al. (1991)] between the Bam HI and Hind III cloning sites.

- 5 Fig. 19 shows the amino acid sequence (one letter code) of the FX-mtlec part of the H6FX-mtlec fusion protein produced by pT7H6FX-mtlec.

Fig. 20 shows an outline of the pT7H6FX-mtCTLD expression plasmid. The FX-mtCTLD fragment was inserted into pT7H6
10 [Christensen et al. (1991)] between the Bam HI and Hind III cloning sites.

Fig. 21 shows the amino acid sequence (one letter code) of the FX-mtCTLD part of the H6FX-mtCTLD fusion protein produced by pT7H6FX-mtCTLD.

- 15 Fig. 22 shows an outline of the pPmtlec phagemid. The Pmtlec fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.

Fig. 23 shows the amino acid sequence (one letter code)
20 of the Pmtlec part of the Pmtlec-gene III fusion protein produced by pPmtlec.

Fig. 24 shows an outline of the pPmtCTLD phagemid. The PmtCTLD fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) be-
25 tween the Sfi I and Not I restriction sites.

Fig. 25 shows the amino acid sequence (one letter code) of the PmtCTLD part of the PmtCTLD-gene III fusion protein produced by pPmtCTLD.

Fig. 26 shows an ELISA-type analysis of Phtlec-, PhTN3-, and M13K07 helper phage binding to anti-tetranectin or BSA. Panel A: Analysis with 3% skimmed milk/5 mM EDTA as blocking reagent. Panel B: Analysis with 3% skimmed milk
5 as blocking reagent.

Fig. 27 shows an ELISA-type analysis of Phtlec-, PhTN3-, and M13K07 helper phage binding to plasminogen (Plg) and BSA. Panel A: Analysis with 3% skimmed milk/5 mM EDTA as blocking reagent. Panel B: Analysis with 3% skimmed milk
10 as blocking reagent.

Fig. 28 shows an ELISA-type analysis of the B series and C series polyclonal populations, from selection round 2, binding to plasminogen (Plg) compared to background.

Fig. 29 Phages from twelve clones isolated from the third
15 round of selection analysed for binding to hen egg white lysozyme, human β_2 -microglobulin and background in an ELISA-type assay.

Fig. 30 shows the amino acid sequence (one letter code) of the PrMBP part of the PrMBP-gene III fusion protein
20 produced by pPrMBP.

Fig. 31 shows an outline of the pPrMBP phagemid. The PrMBP fragment was inserted into the phagemid pCANTAB 5E (Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.

25 Fig. 32 shows the amino acid sequence (one letter code) of the PhSP-D part of the PhSP-D-gene III fusion protein produced by pPhSP-D.

Fig. 33 shows an outline of the pPhSP-D phagemid. The PhSP-D fragment was inserted into the phagemid pCANTAB 5E

(Amersham Pharmacia Biotech, code no. 27-9401-01) between the Sfi I and Not I restriction sites.

Fig. 34. Phages from 48 clones isolated from the third round of selection in the #1 series analysed for binding to hen egg white lysozyme and to A-HA in an ELISA-type assay.

Fig. 35. Phages from 48 clones isolated from the third round of selection in the #4 series analysed for binding to hen egg white lysozyme and to A-HA in an ELISA-type assay.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

The terms "C-type lectin-like protein" and "C-type lectin" are used to refer to any protein present in, or encoded in the genomes of, any eukaryotic species, which protein contains one or more CTLDs or one or more domains belonging to a subgroup of CTLDs, the CRDs, which bind carbohydrate ligands. The definition specifically includes membrane attached C-type lectin-like proteins and C-type lectins, "soluble" C-type lectin-like proteins and C-type lectins lacking a functional transmembrane domain and variant C-type lectin-like proteins and C-type lectins in which one or more amino acid residues have been altered in vivo by glycosylation or any other post-synthetic modification, as well as any product that is obtained by chemical modification of C-type lectin-like proteins and C-type lectins.

In the claims and throughout the specification certain alterations may be defined with reference to amino acid residue numbers of a CTLD domain or a CTLD-containing

protein. The amino acid numbering starts at the first N-terminal amino acid of the CTLD or the native or artificial CTLD-containing protein product, as the case may be, which shall in each case be indicated by unambiguous external literature reference or internal reference to a figure contained herein within the textual context.

The terms "amino acid", "amino acids" and "amino acid residues" refer to all naturally occurring L- α -amino acids. This definition is meant to include norleucine, ornithine, and homocysteine. The amino acids are identified by either the single-letter or three-letter designations:

	Asp D aspartic acid	Ile I isoleucine
	Thr T threonine	Leu L leucine
	Ser S serine	Tyr Y tyrosine
15	Glu E glutamic acid	Phe F phenylalanine
	Pro P proline	His H histidine
	Gly G glycine	Lys K lysine
	Ala A alanine	Arg R arginine
	Cys C cysteine	Trp W tryptophan
20	Val V valine	Gln Q glutamine
	Met M methionine	Asn N asparagine
	Nle J norleucine	Orn O ornithine
	Hcy U homocysteine	Xxx X any L- α -amino acid.

The naturally occurring L- α -amino acids may be classified according to the chemical composition and properties of their side chains. They are broadly classified into two groups, charged and uncharged. Each of these groups is divided into subgroups to classify the amino acids more accurately:

30 A. Charged Amino Acids

Acidic Residues:	Asp, Glu
Basic Residues:	Lys, Arg, His, Orn

B. Uncharged Amino Acids

Hydrophilic Residues: Ser, Thr, Asn, Gln

Aliphatic Residues: Gly, Ala, Val, Leu, Ile,
Nle

5 Non-polar Residues: Cys, Met, Pro, Hcy

Aromatic Residues: Phe, Tyr, Trp

The terms "amino acid alteration" and "alteration" refer to amino acid substitutions, deletions or insertions or any combinations thereof in a CTLD amino acid sequence.

10 In the CTLD variants of the present invention such alteration is at a site or sites of a CTLD amino acid sequence. Substitutional variants herein are those that have at least one amino acid residue in a native CTLD sequence removed and a different amino acid inserted in its

15 place at the same position. The substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule.

The designation of the substitution variants herein consists of a letter followed by a number followed by a letter.

20 The first (leftmost) letter designates the amino acid in the native (unaltered) CTLD or CTLD-containing protein. The number refers to the amino acid position where the amino acid substitution is being made, and the

25 second (righthand) letter designates the amino acid that is used to replace the native amino acid. As mentioned above, the numbering starts with "1" designating the N-terminal amino acid sequence of the CTLD or the CTLD-containing protein, as the case may be. Multiple alterations

30 are separated by a comma (,) in the notation for ease of reading them.

The terms "nucleic acid molecule encoding", "DNA sequence encoding", and "DNA encoding" refer to the order or se-

quence of deoxyribonucleotides along a strand of deoxy-
ribonucleic acid. The order of these deoxyribonucleotides
determines the order of amino acids along the polypeptide
chain. The DNA sequence thus encodes the amino acid se-
5 quence.

The terms "mutationally randomised sequence", "randomised
polypeptide segment", "randomised amino acid sequence",
"randomised oligonucleotide" and "mutationally randomised
sequence", as well as any similar terms used in any con-
10 text to refer to randomised sequences, polypeptides or
nucleic acids, refer to ensembles of polypeptide or nu-
cleic acid sequences or segments, in which the amino acid
residue or nucleotide at one or more sequence positions
may differ between different members of the ensemble of
15 polypeptides or nucleic acids, such that the amino acid
residue or nucleotide occurring at each such sequence po-
sition may belong to a set of amino acid residues or nu-
cleotides that may include all possible amino acid resi-
dues or nucleotides or any restricted subset thereof.
20 Said terms are often used to refer to ensembles in which
the number of amino acid residues or nucleotides is the
same for each member of the ensemble, but may also be
used to refer to such ensembles in which the number of
amino acid residues or nucleotides in each member of the
25 ensemble may be any integer number within an appropriate
range of integer numbers.

II. Construction and utility of combinatorial CTLD li- braries

Several systems displaying phenotype, in terms of puta-
30 tive ligand binding modules or modules with putative en-
zymatic activity, have been described. These include:
phage display (e.g. the filamentous phage fd [Dunn
(1996), Griffiths and Duncan (1998), Marks et al.

(1992)], phage lambda [Mikawa et al. (1996)], display on eukarotic virus (e.g. baculovirus [Ernst et al. (2000)]), cell display (e.g. display on bacterial cells [Benhar et al. (2000)], yeast cells [Boder and Wittrup (1997)], and
5 mammalian cells [Whitehorn et al. (1995)], ribosome linked display [Schaffitzel et al. (1999)], and plasmid linked display [Gates et al. (1996)].

The most commonly used method for phenotype display and linking this to genotype is by phage display. This is ac-
10 complished by insertion of the reading frame encoding the scaffold protein or protein of interest into an intra-domain segment of a surface exposed phage protein. The filamentous phage fd (e.g. M13) has proven most useful for this purpose. Polypeptides, protein domains, or pro-
15 teins are the most frequently inserted either between the "export" signal and domain 1 of the fd gene III protein or into a so-called hinge region between domain 2 and domain 3 of the fd-phage gene III protein. Human antibodies are the most frequently used proteins for the isolation
20 of new binding units, but other proteins and domains have also been used (e.g. human growth hormone [Bass et al. (1990)], alkaline phosphatase [McCafferty et al. (1991)], β -lactamase inhibitory protein [Huang et al. (2000)], and cytotoxic T lymphocyte-associated antigen 4 [Hufton et
25 al. (2000)]. The antibodies are often expressed and presented as scFv or Fab fusion proteins. Three strategies have been employed. Either a specific antibody is used as a scaffold for generating a library of mutationally randomised sequences within the antigen binding clefts [e.g.
30 Fuji et al. (1998)] or libraries representing large ensembles of human antibody encoding genes from non-immunised hosts [e.g. Nissim et al. (1994)] or from immunised hosts [e.g. Cyr and Hudspeth (2000)] are cloned into the fd phage vector.

The general procedure for accomplishing the generation of a display system for the generation of CTLD libraries comprise essentially

- 5 (1) identification of the location of the loop-region, by referring to the 3D structure of the CTLD of choice, if such information is available, or, if not, identification of the sequence locations of the $\beta 2$ -, $\beta 3$ - and $\beta 4$ strands by sequence alignment with the sequences shown in Fig. 1, as
10 aided by the further corroboration by identification of sequence elements corresponding to the $\beta 2$ and $\beta 3$ consensus sequence elements and $\beta 4$ -strand characteristics, also disclosed above;
- 15 (2) subcloning of a nucleic acid fragment encoding the CTLD of choice in a protein display vector system with or without prior insertion of endonuclease restriction sites close to the sequences encoding $\beta 2$, $\beta 3$ and $\beta 4$; and
- 20 (3) substituting the nucleic acid fragment encoding some or all of the loop-region of the CTLD of choice with randomly selected members of an ensemble consisting of a multitude of nucleic acid fragments which after insertion into the nucleic acid context encoding the receiving framework
25 will substitute the nucleic acid fragment encoding the original loop-region polypeptide fragments with randomly selected nucleic acid fragments. Each of the cloned nucleic acid fragments, encoding a new polypeptide replacing an original
30 loop-segment or the entire loop-region, will be decoded in the reading frame determined within its new sequence context.

Nucleic acid fragments may be inserted in specific locations into receiving nucleic acids by any common method of molecular cloning of nucleic acids, such as by appropriately designed PCR manipulations in which chemically synthesized nucleic acids are copy-edited into the receiving nucleic acid, in which case no endonuclease restriction sites are required for insertion. Alternatively, the insertion/excision of nucleic acid fragments may be facilitated by engineering appropriate combinations of endonuclease restriction sites into the target nucleic acid into which suitably designed oligonucleotide fragments may be inserted using standard methods of molecular cloning of nucleic acids.

It will be apparent that interesting CTLD variants isolated from CTLD libraries in which restriction endonuclease sites have been inserted for convenience may contain mutated or additional amino acid residues that neither correspond to residues present in the original CTLD nor are important for maintaining the interesting new affinity of the CTLD variant. If desirable, e.g. in case the product needs to be rendered as non-immunogenic as possible, such residues may be altered or removed by back-mutation or deletion in the specific clone, as appropriate.

The ensemble consisting of a multitude of nucleic acid fragments may be obtained by ordinary methods for chemical synthesis of nucleic acids by directing the step-wise synthesis to add pre-defined combinations of pure nucleotide monomers or a mixture of any combination of nucleotide monomers at each step in the chemical synthesis of the nucleic acid fragment. In this way it is possible to generate any level of sequence degeneracy, from one unique nucleic acid sequence to the most complex mixture,

which will represent a complete or incomplete representation of maximum number unique sequences of 4^N , where N is the number of nucleotides in the sequence.

Complex ensembles consisting of multitudes of nucleic acid fragments may, alternatively, be prepared by generating mixtures of nucleic acid fragments by chemical, physical or enzymatic fragmentation of high-molecular mass nucleic acid compositions like, e.g., genomic nucleic acids extracted from any organism. To render such mixtures of nucleic acid fragments useful in the generation of molecular ensembles, as described here, the crude mixtures of fragments, obtained in the initial cleavage step, would typically be size-fractionated to obtain fragments of an approximate molecular mass range which would then typically be adjoined to a suitable pair of linker nucleic acids, designed to facilitate insertion of the linker-embedded mixtures of size-restricted oligonucleotide fragments into the receiving nucleic acid vector.

To facilitate the construction of combinatorial CTLD libraries in tetranectin, the model CTLD of the preferred embodiment of the invention, suitable restriction sites located in the vicinity of the nucleic acid sequences encoding $\beta 2$, $\beta 3$ and $\beta 4$ in both human and murine tetranectin were designed with minimal perturbation of the polypeptide sequence encoded by the altered sequences. It was found possible to establish a design strategy, as detailed below, by which identical endonuclease restriction sites could be introduced at corresponding locations in the two sequences, allowing interesting loop-region variants to be readily excised from a recombinant murine CTLD and inserted correctly into the CTLD framework of human tetranectin or vice versa.

- Analysis of the nucleotide sequence encoding the mature form of human tetranectin reveals (Fig. 2) that a recognition site for the restriction endonuclease *Bgl* II is found at position 326 to 331 (AGATCT), involving the encoded residues Glu109, Ile110, and Trp111 of β 2, and that a recognition site for the restriction endonuclease *Kas* I is found at position 382 to 387 (GGCGCC), involving the encoded amino acid residues Gly128 and Ala129 (located C-terminally in loop 2).
- 10 Mutation, by site directed mutagenesis, of G513 to A and of C514 to T in the nucleotide sequence encoding human tetranectin would introduce a *Mun* I restriction endonuclease recognition site therein, located at position 511 to 516, and mutation of G513 to A in the nucleotide sequence encoding murine tetranectin would introduce a *Mun* I restriction endonuclease site therein at a position corresponding to the *Mun* I site in human tetranectin, without affecting the amino acid sequence of either of the encoded protomers. Mutation, by site directed
- 15 mutation, of C327 to G and of G386 to C in the nucleotide sequence encoding murine tetranectin would introduce a *Bgl* II and a *Kas* I restriction endonuclease recognition site, respectively, therein. Additionally, A325 in the nucleotide sequence encoding murine tetranectin is
- 20 mutagenized to a G. These three mutations would affect the encoded amino acid sequence by substitution of Asn109 to Glu and Gly129 to Ala, respectively. Now, the restriction endonuclease *Kas* I is known to exhibit marked site preference and cleaves only slowly the tetranectin coding region. Therefore, a recognition site for another restriction endonuclease substituting the *Kas* I site is
- 25 preferred (e.g. the recognition site for the restriction endonuclease *Kpn* I, recognition sequence GGTACC). The nucleotide and amino acid sequences of the resulting
- 30

tetranectin derivatives, human tetranectin lectin (htlec) and murine tetranectin lectin (mtlec) are shown in Fig. 3. The nucleotide sequences encoding the htlec and mtlec protomers may readily be subcloned into devices enabling protein display of the linked nucleotide sequence (e.g. phagemid vectors) and into plasmids designed for heterologous expression of protein [e.g. pT7H6, Christensen et al. (1991)]. Other derivatives encoding only the mutated CTLDs of either htlec or mtlec (htCTLD and mtCTLD, respectively) have also been constructed and subcloned into phagemid vectors and expression plasmids, and the nucleotide and amino acid sequences of these CTLD derivatives are shown in Fig. 4.

The presence of a common set of recognition sites for the restriction endonucleases *Bgl* II, *Kas* I or *Kpn* I, and *Mun* I in the ensemble of tetranectin and CTLD derivatives allows for the generation of protein libraries with randomised amino acid sequence in one or more of the loops and at single residue positions in β 4 comprising the lectin ligand binding region by ligation of randomised oligonucleotides into properly restricted phagemid vectors encoding htlec, mtlec, htCTLD, or mtCTLD derivatives.

After rounds of selection on specific targets (e.g. eukaryotic cells, virus, bacteria, specific proteins, polysaccharides, other polymers, organic compounds etc.) DNA may be isolated from the specific phages, and the nucleotide sequence of the segments encoding the ligand-binding region determined, excised from the phagemid DNA and transferred to the appropriate derivative expression vector for heterologous production of the desired product. Heterologous production in a prokaryote may be preferred because an efficient protocol for the isolation and refolding of tetranectin and derivatives has been reported

(International Patent Application Publication WO 94/18227 A2).

A particular advantage gained by implementing the technology of the invention, using tetranectin as the scaffold structure, is that the structures of the murine and human tetranectin scaffolds are almost identical, allowing loop regions to be swapped freely between murine and human tetranectin derivatives with retention of functionality. Swapping of loop regions between the murine and the human framework is readily accomplished within the described system of tetranectin derivative vectors, and it is anticipated, that the system can be extended to include other species (e.g. rat, old and new world monkeys, dog, cattle, sheep, goat etc.) of relevance in medicine or veterinary medicine in view of the high level of homology between man and mouse sequences, even at the genetic level. Extension of this strategy to include more species may be rendered possible as and when tetranectin is eventually cloned and/or sequenced from such species.

Because the C-type lectin ligand-binding region represents a different topological unit compared to the antigen binding clefts of the antibodies, we envisage that the selected binding specificities will be of a different nature compared to the antibodies. Further, we envisage that the tetranectin derivatives may have advantages compared to antibodies with respect to specificity in binding sugar moieties or polysaccharides. The tetranectin derivatives may also be advantageous in selecting binding specificities against certain natural or synthetic organic compounds.

Several CTLDs are known to bind calcium ions, and binding of other ligands is often either dependent on calcium (e.g. the collectin family of C-type lectins, where the

calcium ion bound in site 2 is directly involved in binding the sugar ligand [Weis and Drickamer (1996)] or sensitive to calcium (e.g. tetranectin, where binding of calcium involves more of the side chains known otherwise to be involved in plasminogen kringle 4 binding [Grav-
5 ersen et al. (1998)]). The calcium binding sites characteristic of the C-type lectin-like protein family are comprised by residues located in loop 1, loop 4 and β -strand 4 and are dependent on the presence of a proline
10 residue (often interspacing loop 3 and loop 4 in the structure), which upon binding is found invariantly in the *cis* conformation. Moreover, binding of calcium is known to enforce structural changes in the CTLD loop-region [Ng et al. (1998a,b)]. We therefore envisage, that
15 binding to a specific target ligand by members of combinatorial libraries with preserved CTLD metal binding sites may be modulated by addition or removal of divalent metal ions (e.g. calcium ions) either because the metal ion may be directly involved in binding, because it is a
20 competitive ligand, or because binding of the metal ion enforces structural rearrangements within the putative binding site.

The trimeric nature of several members of the C-type lectin and C-type lectin-like protein family, including
25 tetranectin, and the accompanying avidity in binding may also be exploited in the creation of binding units with very high binding affinity.

As can be appreciated from the disclosure above, the present invention has a broad general scope and a wide area
30 of application. Accordingly, the following examples, describing various embodiments thereof, are offered by way

of illustration only, not by way of limitation.

Example 1

5 Construction of tetranectin derived *E.coli* expression plasmids and phagemids

The expression plasmid pT7H6FX-htlec, encoding the FX-htlec (SEQ ID NO:01) part of full length H6FX-htlec fusion protein, was constructed by a series of four consecutive site-directed mutagenesis experiments starting
10 from the expression plasmid pT7H6-rTN 123 [Holtet et al. (1997)] using the QuickChange™ Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA) and performed as described by the manufacturer. Mismatching primer pairs introducing the desired mutations were supplied by DNA Technology
15 (Aarhus, Denmark). An outline of the resulting pT7H6FX-htlec expression plasmid is shown in Fig. 5, and the nucleotide sequence of the FX-htlec encoding insert is given as SEQ ID NO:01. The amino acid sequence of the FX-htlec part of the H6FX-htlec fusion protein is shown in
20 Fig. 6 and given as SEQ ID NO:02.

The expression plasmid pT7H6FX-htCTLD, encoding the FX-htCTLD (SEQ ID NO: 03) part of the H6FX-htCTLD fusion protein, was constructed by amplification and subcloning into the plasmid pT7H6 (i.e. amplification in a polymerase chain reaction using the expression plasmid pT7H6-htlec as template, and otherwise the primers, conditions, and subcloning procedure described for the construction of the expression plasmid pT7H6TN3 [Holtet et al. (1997)]). An outline of the resulting pT7H6FX-htCTLD
30 expression plasmid is shown in Fig. 7, and the nucleotide sequence of the FX-htCTLD encoding insert is given as SEQ ID NO:03. The amino acid sequence of the FX-htCTLD part

of the H6FX-htCTLD fusion protein is shown in Fig. 8 and given as SEQ ID NO:04.

The phagemids, pPhTN and pPhTN3, were constructed by ligation of the *Sfi* I and *Not* I restricted DNA fragments amplified from the expression plasmids pT7H6-rTN 123 (with the oligonucleotide primers 5-CGGCTGAGCGGCCCA-GCCGGCCATGGCCGAGCCACCAACCCAGAAGC-3' [SEQ ID NO:05] and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID NO:06]) and pT7H6FX-htCTLD (with the oligonucleotide primers 5'-CGGCTGAGCGGCCCAAGCCGGCCATGGCCGCCCTGCAGACGGTC-3' [SEQ ID NO:07] and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID NO:06]), respectively, into a *Sfi* I and *Not* I precut vector, pCANTAB 5E supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) using standard procedures. Outlines of the resulting pPhTN and pPhTN3 phagemids are shown in Fig. 9 and Fig. 11, respectively, and the nucleotide sequences of the PhTN and PhTN3 inserts are given as SEQ ID NO:08 and SEQ ID NO:10, respectively. The amino acid sequences encoded by the PhTN and PhTN3 inserts are shown in Fig 10 (SEQ ID NO:09) and Fig. 12 (SEQ ID NO:11), respectively.

The phagemids, pPhTlec and pPhtCTLD, were constructed by ligation of the *Sfi* I and *Not* I restricted DNA fragments amplified from the expression plasmids pT7H6FX-htlec (with the oligonucleotide primers 5-CGGCTGAGCGGCCCAAGCCGGCCATGGCCGAGCCACCAACCCAGAAGC-3' [SEQ ID NO:05] and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID NO:06]) and pT7H6FX-htCTLD (with the oligonucleotide primers 5'-CGGCTGAGCGGCCCAAGCCGGCCATGGCCGCCCTGCAGACGGTC-3' [SEQ ID NO:07] and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID NO:06]), respectively, into a *Sfi* I and *Not* I precut vector, pCANTAB 5E supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) using standard procedures. Outlines

of the resulting pPhtlec and pPhtCTLD phagemids are shown in Fig. 13 and Fig. 15, respectively, and the nucleotide sequences of the Phtlec and PhtCTLD inserts are given as SEQ ID NO:12 and SEQ ID NO:14, respectively. The amino acid sequences encoded by the Phtlec and PhtCTLD inserts are shown in Fig. 14 (SEQ ID NO:13) and Fig. 16 (SEQ ID NO:15), respectively.

A plasmid clone, pUC-mtlec, containing the nucleotide sequence corresponding to the murine tetranectin derivative mtlec (Fig. 3 and SEQ ID NO:16) was constructed by four successive subclonings of DNA subfragments in the following way: First, two oligonucleotides 5'-CGGAATTCGAGTCACCCACTCCCAAGGCCAAGAAGGCTGCAAATGCCAAGAAA-GATTTGGTGAGCTCAAAGATGTTC-3' (SEQ ID NO:17) and 5'-GCG-GATCCAGGCCTGCTTCTCCTTCAGCAGGGCCACCTCCTGGGCCAGGACATCCAT-CCTGTTCTTGAGCTCCTCGAACATCTTTGAGCTCACC-3' (SEQ ID NO:18) were annealed and after a filling in reaction cut with the restriction endonucleases *Eco* RI (GAATTC) and *Bam* HI (GGATCC) and ligated into *Eco* RI and *Bam* HI precut pUC18 plasmid DNA. Second, a pair of oligonucleotides 5'-GCA-GGCCTTACAGACTGTGTGCCTGAAGGGCACCAAGGTGAACTTGAAGTGCCTCCT-GGCCTTCACCCAACCGAAGACCTTCCATGAGGCGAGCGAG-3' (SEQ ID NO:19) and 5'-CCGCATGCTTCGAACAGCGCCTCGTTCTCTAGCTCTGAC-TGCGGGGTGCCCAGCGTGCCCCCTTGCGAGATGCAGTCCTCGCTCGCCTCATGG-3' (SEQ ID NO:20) was annealed and after a filling in reaction cut with the restriction endonucleases *Stu* I (AGGCCT) and *Sph* I (GCATGC) and ligated into the *Stu* I and *Sph* I precut plasmid resulting from the first ligation. Third, an oligonucleotide pair 5'-GGTTCGAATACGCGC-GCCACAGCGTGGGCAACGATGCGGAGATCTAAATGCTCCCAATTGC-3' (SEQ ID NO:21) and 5'-CCAAGCTTCACAATGGCAAACCTGGCAGATGTAGGGCAATTGG-GAGCATTTAGATC-3' (SEQ ID NO: 22) was annealed and after a filling in reaction cut with the restriction endonucleases *Bst*B I (TTCGAA) and *Hind* III (AAGCTT) and ligated

into the *Bst*B I and *Hind* III precut plasmid resulting from the second ligation. Fourth, an oligonucleotide pair 5'-CGGAGATCTGGCTGGGCCTCAACGACATGGCCGCGGAAGGCGCCTGGGTGGA-CATGACCGGTACCCTCCTGGCCTACAAGAACTGG-3' (SEQ ID NO:23) and
5 5'-GGGCAATTGATCGCGGCATCGCTTGTCGAACCTCTTGCCGTTGGCTGCGCCAG-ACAGGGCGGCGCAGTTCTCGGCTTTGCCGCCGTCGGGTTGCGTCGTGATCTCCGTC-TCCCAGTTCTTGTAGGCCAGG-3' (SEQ ID NO:24) was annealed and after a filling in reaction cut with the restriction endonucleases *Bgl* II (AGATCT) and *Mun* I (CAATTG) and ligated
10 into the *Bgl* II and *Mun* I precut plasmid resulting from the third ligation. An outline of the pUC-mtlec plasmid is shown in Fig. 17, and the resulting nucleotide sequence of the *Eco* RI to *Hind* III insert is given as SEQ ID NO:16.

15 The expression plasmids pT7H6FX-mtlec and pT7H6FX-mtCTLD may be constructed by ligation of the *Bam* HI and *Hind* III restricted DNA fragments, amplified from the pUC-mtlec plasmid with the oligonucleotide primer pair 5'-CTGGGATCC-ATCCAGGGTCGCGAGTCACCCACTCCCAAGG-3' (SEQ ID NO:25) and 5'-
20 CCGAAGCTTACACAATGGCAAACCTGGC-3' (SEQ ID NO:26), and with the oligonucleotide primer pair 5'-CTGGGATCCATCCAGGGTCGCGCCTTACAGACTGTGGTC-3' (SEQ ID NO:27), and 5'-CCGAAGCTT-ACACAATGGCAAACCTGGC-3' (SEQ ID NO:26), respectively, into *Bam* HI and *Hind* III precut pT7H6 vector using standard
25 procedures. An outline of the expression plasmids pT7H6FX-mtlec and pT7H6FX-mtCTLD is shown in Fig. 18 and Fig. 20, respectively, and the nucleotide sequences of the FX-mtlec and FX-mtCTLD inserts are given as SEQ ID NO:28 and SEQ ID NO:30, respectively. The amino acid sequences of the FX-mtlec and FX-mtCTLD parts of the fusion
30 proteins H6FX-mtlec and H6FX-mtCTLD fusion proteins are shown in Fig. 19 (SEQ ID NO:29) and Fig. 21 (SEQ ID NO:31), respectively.

The phagemids pPmtlec and pPmtCTLD may be constructed by ligation of the Sfi I and Not I restricted DNA fragments (amplified from the pUC-mtlec plasmid with the oligonucleotide primer pair 5-

5 CGGCTGAGCGGCCCAGCCGGCCATGGCCGAGTCACCCACTCCCAAGG-3' [SEQ ID NO:32], and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID NO:33] and with the oligonucleotide primers 5'-

CGGCTGAGCGGCCCAGCCGGCCATGGCCGCTTACAGACTGTGGTC-3' [SEQ ID NO:34] and 5'-CCTGCGGCCGCCACGATCCCGAACTGG-3' [SEQ ID

10 NO:33], respectively) into a Sfi I and Not I precut vector pCANTAB 5E supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) using standard procedures. Outlines of the pPmtlec and pPmtCTLD plasmids are shown in Fig. 22 and Fig. 24, respectively, and the resulting nucleotide

15 sequences of the Pmtlec and PmtCTLD inserts are given as SEQ ID NO:35 and SEQ ID NO:37, respectively. The amino acid sequences encoded by the Pmtlec and PmtCTLD inserts are shown in Fig. 23 (SEQ ID NO: 36) and Fig. 25 (SEQ ID NO: 38), respectively.

20

20 Example 2

Demonstration of successful display of Phtlec and PhTN3 on phages.

In order to verify that the Phtlec and PhTN3 Gene III fusion proteins can indeed be displayed by the recombinant

25 phage particles, the phagemids pPhtlec and pPhTN3 (described in Example 1) were transformed into *E. coli* TG1 cells and recombinant phages produced upon infection with the helper phage M13KO7. Recombinant phages were isolated by precipitation with poly(ethylene glycol) (PEG 8000)

30 and samples of Phtlec and PhTN3 phage preparations as well as a sample of helper phage were subjected to an ELISA-type sandwich assay, in which wells of a Maxisorb (Nunc) multiwell plate were first incubated with anti-

human tetranectin or bovine serum albumin (BSA) and blocked in skimmed milk or skimmed milk/EDTA. Briefly, cultures of pPhtlec and pPhTN3 phagemid transformed TG1 cells were grown at 37 °C in 2xTY-medium supplemented with 2% glucose and 100 mg/L ampicillin until A_{600} reached 0.5. By then the helper phage, M13K07, was added to a concentration of 5×10^9 pfu/mL. The cultures were incubated at 37 °C for another 30 min before cells were harvested by centrifugation and resuspended in the same culture volume of 2xTY medium supplemented with 50 mg/L kanamycin and 100 mg/L ampicillin and transferred to a fresh set of flasks and grown for 16 hours at 25 °C. Cells were removed by centrifugation and the phages precipitated from 20 mL culture supernatant by the addition of 6 mL of ice cold 20% PEG 8000, 2.5 M NaCl. After mixing the solution was left on ice for one hour and centrifuged at 4 °C to isolate the precipitated phages. Each phage pellet was resuspended in 1 mL of 10 mM tris-HCl pH 8, 1 mM EDTA (TE) and incubated for 30 min before centrifugation. The phage containing supernatant was transferred to a fresh tube. Along with the preparation of phage samples, the wells of a Maxisorb plate was coated overnight with (70 μ L) rabbit anti-human tetranectin (a polyclonal antibody from DAKO A/S, code no. A0371) in a 1:2000 dilution or with (70 μ L) BSA (10 mg/mL). Upon coating, the wells were washed three times with PBS (2.68 mM KCl, 1.47 mM KH_2PO_4 , 137 mM NaCl, 8.10 mM Na_2HPO_4 , pH 7.4) and blocked for one hour at 37 °C with 280 μ L of either 3% skimmed milk in PBS, or 3% skimmed milk, 5mM EDTA in PBS. Anti-tetranectin coated and BSA coated wells were then incubated with human Phtlec-, PhTN3-, or helper phage samples for 1 hour and then washed 3 times in PBS buffer supplemented with the appropriate blocking agent. Phages in the wells were detected after incubation with HRP-conjugated anti-phage conjugate (Amersham Pharmacia,

code no. 27-9421-01) followed by further washing. HRP activities were then measured in a 96-well ELISA reader using a standard HRP chromogenic substrate assay.

5 Phtlec and PhTN3 phages produced strong responses (14 times background) in the assay, irrespective of the presence or absence of EDTA in the blocking agent, whereas helper phage produced no response above background readings in either blocking agent. Only low binding to BSA was observed (Fig. 26).

10 It can therefore be concluded that the human Phtlec and PhTN3 phages both display epitopes that are specifically recognized by the anti-human tetranectin antibody.

Example 3

15 Demonstration of authentic ligand binding properties of Phtlec and PhTN3 displayed on phage

The apo-form of the CTLD domain of human tetranectin binds in a lysine-sensitive manner specifically to the kringle 4 domain of human plasminogen [Graversen et al. 20 (1998)]. Binding of tetranectin to plasminogen can be inhibited by calcium which binds to two sites in the ligand-binding site in the CTLD domain (K_d approx. 0.2 millimolar) or by lysine-analogues like AMCHA (6-amino-cyclohexanoic acid), which bind specifically in the two 25 stronger lysine-binding sites in plasminogen of which one is located in kringle 1 and one is located in kringle 4 (K_d approx. 15 micromolar).

To demonstrate specific AMCHA-sensitive binding of human Phtlec and PhTN3 phages to human plasminogen, an ELISA 30 assay, in outline similar to that employed to demonstrate

the presence of displayed Phlec and PhCTLD GIII fusion proteins on the phage particles (cf. Example 2), was devised.

5 Wells were coated with solutions of human plasminogen (10 $\mu\text{g/mL}$), with or without addition of 5mM AMCHA. Control wells were coated with BSA. Two identical arrays were established, one was subjected to blocking of excess binding capacity with 3% skimmed milk, and one was blocked using 3% skimmed milk supplemented with 5mM EDTA. Where
10 appropriate, blocking, washing and phage stock solutions were supplemented by 5mM AMCHA. The two arrays of wells were incubated with either Phlec-, or PhTN3-, or helper phage samples, and after washing the amount of phage bound in each well was measured using the HRP-conjugated
15 antiphage antibody as above. The results are shown in Fig. 27, panels A and B, and can be summarized as follows

- 20 (a) In the absence of AMCHA, binding of human Phlec phages to plasminogen-coated wells generated responses at 8-10 times background levels using either formulation of blocking agent, whereas human PhTN3 phages generated responses at 4 (absence of EDTA) or 7 (presence of EDTA) times background response levels.
- 25 (b) In the presence of 5mM AMCHA, binding of human Phlec- and PhTN3 phages to plasminogen was found to be completely abolished.
- (c) Phlec and PhTN3 phages showed no binding to BSA, and control helper phages showed no binding to any of the immobilized substances.
- 30 (d) Specific binding of human Phlec and PhTN3 phages to a specific ligand at moderate binding strength

(about 20 micromolar level) can be detected with high efficiency at virtually no background using a skimmed-milk blocking agent, well-known in the art of combinatorial phage technology as a preferred agent effecting the reduction of non-specific binding.

In conclusion, the results show that the Phtlec and PhTN3 Gene III fusion proteins displayed on the phage particles exhibit plasminogen-binding properties corresponding to those of authentic tetranectin, and that the physical and biochemical properties of Phtlec and PhTN3 phages are compatible with their proposed use as vehicles for the generation of combinatorial libraries from which CTLD derived units with new binding properties can be selected.

Example 4

Construction of the phage libraries Phtlec-lb001 and Phtlec-lb002.

All oligonucleotides used in this example were supplied by DNA Technology (Aarhus, Denmark).

The phage library Phtlec-lb001, containing random amino acid residues corresponding to Phtlec (SEQ ID NO: 12) positions 141-146 (loop 3), 150-153 (part of loop 4), and residue 168 (Phe in β 4), was constructed by ligation of 20 μ g *Kpn*I and *Mun*I restricted pPhtlec phagemid DNA (cf, Example 1) with 10 μ g of *Kpn*I and *Mun*I restricted DNA fragment amplified from the oligonucleotide htlec-lib1-tp (SEQ ID NO: 39), where N denotes a mixture of 25% of each of the nucleotides T, C, G, and A, respectively and S denotes a mixture of 50 % of C and G, encoding the appropriately randomized nucleotide sequence and the oligonu-

cleotides htlec-lib1-rev (SEQ ID NO: 40) and htlec-lib1/2-fo (SEQ ID NO: 41) as primers using standard conditions. The ligation mixture was used to transform so-called electrocompetent *E. coli* TG-1 cells by electroporation using standard procedures. After transformation
5 the *E. coli* TG-1 cells were plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated over night at 30°C.

The phage library Phtlec-lb002, containing random amino
10 acid residues corresponding to Phtlec (SEQ ID NO: 12) positions 121-123, 125 and 126 (most of loop 1), and residues 150-153 (part of loop 4) was constructed by ligation of 20 µg *Bgl*III and *Mun*I restricted pPhtlec phagemid DNA (cf, EXAMPLE 1) with 15 µg of *Bgl*III and *Mun*I restricted
15 DNA fragment amplified from the pair of oligonucleotides htlec-lib2-tprev (SEQ ID NO: 42) and htlec-lib2-tpfo (SEQ ID NO: 43), where N denotes a mixture of 25% of each of the nucleotides T, C, G, and A, respectively and S denotes a mixture of 50 % of C and G, encoding the appropriately randomized nucleotide sequence and the oligonucleotides htlec-lib2-rev (SEQ ID NO: 44) and htlec-lib1/2-fo (SEQ ID NO: 41) as primers using standard conditions. The ligation mixture was used to transform so-called electrocompetent *E. coli* TG-1 cells by electroporation using standard procedures. After transformation
20 the *E. coli* TG-1 cells were plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated overnight at 30°C.
25

The titer of the libraries Phtlec-lb001 and -lb002 was
30 determined to $1.4 \cdot 10^9$ and $3.2 \cdot 10^9$ clones, respectively. Six clones from each library were grown and phagemid DNA isolated using a standard miniprep procedure, and the nucleotide sequence of the loop-region determined (DNA

Technology, Aarhus, Denmark). One clone from each library failed, for technical reasons, to give reliable nucleotide sequence, and one clone from Phtlec-lib001 apparently contained a major deletion. The variation of nucleotide sequences, compared to Phtlec (SEQ ID NO: 12),
5 of the loop-regions of the other nine clones (lb001-1, lb001-2, lb001-3, lb001-4, lb002-1, lb002-2, lb002-3, lb002-4, and lb002-5) is shown in Table 3.

Table 3: Variation of Phtlec loop derivatives isolated from the libraries Phtlec-lb001 and -lb002. ($\beta 2$ and $\beta 3$ consensus elements are indicated)

[illegible]

Example 5

Construction of the phage library PhtCTLD-lb003

All oligonucleotides used in this example were supplied by DNA Technology (Aarhus, Denmark).

- 5 The phage library PhtCTLD-lb003, containing random amino acid residues corresponding to PhtCTLD (SEQ ID NO: 15) positions 77 to 79 and 81 to 82 (loop 1) and 108 to 109 (loop 4) was constructed by ligation of 20 µg *Bgl*III and *Mun*I restricted pPhtCTLD phagemid DNA (cf. Example 1) with 10 µg
10 of a *Bgl*III and *Mun*I restricted DNA fragment population encoding the appropriately randomised loop 1 and 4 regions with or without two and three random residue insertions in loop 1 and with three and four random residue insertions in loop 4. The DNA fragment population was amplified, from six
15 so-called assembly reactions combining each of the three loop 1 DNA fragments with each of the two loop 4 DNA fragments as templates and the oligonucleotides TN-lib3-rev (SEQ ID NO: 45) and loop 3-4-5 tagfo (SEQ ID NO: 46) as primers using standard procedures. Each of the three loop 1
20 fragments was amplified in a reaction with either the oligonucleotides loop1b (SEQ ID NO: 47), loop1c (SEQ ID NO: 48), or loop1d (SEQ ID NO: 49) as template and the oligonucleotides TN-lib3-rev (SEQ ID NO: 45) and TN-KpnI-fo (SEQ ID NO: 50) as primers, and each of the two DNA loop 4 frag-
25 ments was amplified in a reaction with either the oligonucleotide loop4b (SEQ ID NO: 51) or loop4c (SEQ ID NO: 52) as template and the oligonucleotides loop3-4rev (SEQ ID NO: 53) and loop3-4fo (SEQ ID NO: 54) as primers using standard procedures. In the oligonucleotide sequences N denotes a
30 mixture of 25% of each of the nucleotides T, C, G, and A, respectively and S denotes a mixture of 50 % of C and G, encoding the appropriately randomized nucleotide sequence. The ligation mixture was used to transform so-called elec-

trocompetent *E. coli* TG-1 cells by electroporation using standard procedures. After transformation the *E. coli* TG-1 cells were plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated over night at 30 °C.

The size of the resulting library, PhtCTLD-lb003, was determined to 1.4×10^{10} clones. Twenty four clones from the library were grown and phages and phagemid DNA isolated. The nucleotide sequences of the loop-regions were determined (DNA Technology, Aarhus, Denmark) and binding to a polyclonal antibody against tetranectin, anti-TN (DAKO A/S, Denmark), analysed in an ELISA-type assay using HRP conjugated anti-gene VIII (Amersham Pharmacia Biotech) as secondary antibody using standard procedures. Eighteen clones were found to contain correct loop inserts, one clone contained the wild type loop region sequence, one a major deletion, two contained two or more sequences, and two clones contained a frameshift mutation in the region. Thirteen of the 18 clones with correct loop inserts, the wild type clone, and one of the mixed isolates reacted strongly with the polyclonal anti-TN antibody. Three of the 18 correct clones reacted weakly with the antibody, whereas, two of the correct clones, the deletion mutant, one of the mixed, and the two frameshift mutants did not show a signal above background.

Example 6

Phage selection by biopanning on anti-TN antibody.

Approximately 10^{11} phages from the PhtCTLD-lb003 library was used for selection in two rounds on the polyclonal anti-TN antibody by panning in Maxisorb immunotubes (NUNC, Denmark) using standard procedures. Fifteen clones out of 7×10^7 from the plating after the second selection round

were grown and phagemid DNA isolated and the nucleotide sequence determined. All 15 clones were found to encode correct and different loop sequences.

5 Example 7

Model selection of CTLD-phages on plasminogen.

I: elution by trypsin digestion after panning.

In order to demonstrate that tetranectin derived CTLD bearing phages can be selected from a population of phages, mixtures of PhtCTLD phages isolated from a *E. coli* TG1 culture transformed with the phagemid pPhtCTLD (cf, EXAMPLE 1) after infection with M13K07 helper phage and phages isolated from a culture transformed with the phagemid pPhtCPB after infection with M13K07 helper phage at ratios of 1:10 and 1:10⁵, respectively were used in a selection experiment using panning in 96-well Maxisorb micro-titerplates (NUNC, Denmark) and with human plasminogen as antigen. The pPhtCPB phagemid was constructed by ligation of the double stranded oligonucleotide (SEQ ID NO: 55) with the appropriate restriction enzyme overhang sequences into KpnI and MunI restricted pPhtCTLD phagemid DNA. The pPhtCPB phages derived upon infection with the helper phages displays only the wild type M13 gene III protein because of the translation termination codons introduced into the CTLD coding region of the resulting pPhtCPB phagemid (SEQ ID NO: 56).

The selection experiments were performed in 96 well micro titer plates using standard procedures. Briefly, in each well 3 µg of human plasminogen in 100 µL PBS (PBS, 0.2 g KCl, 0.2 g KH₂PO₄, 8 g NaCl, 1.44 g Na₂HPO₄, 2H₂O, water to 1 L, and adjusted to pH 7.4 with NaOH) or 100 µL PBS (for analysis of non specific binding) was used for over night coating at 4 °C and at 37 °C for one hour. After washing once with PBS, wells were blocked with 400 µL PBS and 3%

non fat dried milk for one hour at 37°C. After blocking wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of phages suspended in 100 µL PBS, 3% non fat dried milk. The phages were allowed to bind at 37 °C for one hour before washing three times with PBS, Tween 20 and three times with PBS. Bound phages were eluted from each well by trypsin digestion in 100 µL (1 mg/mL trypsin in PBS) for 30 min. at room temperature, and used for infection of exponentially growing *E. coli* TG1 cells before plating and titration on 2xTY agar plates containing 2% glucose and 0.1 mg/mL ampicillin.

Initially (round 1), 10^{12} PhtCTLD phages (A series), a mixture of 10^{10} PhtCTLD phages and 10^{11} PhtCPB phages (B series), or a mixture of 10^6 PhtCTLD and 10^{11} PhtCPB phages (C series) were used. In the following round (round 2) 10^{11} phages of the output from each series were used. Results from the two rounds of selection are summarised in Table 4.

Table 4: Selection of mixtures of PhtCTLD and PhtCPB by panning and elution with trypsin.

		Plasminogen (* 10^5 colonies)	Blank (* 10^5 colonies)
Round 1	A	113.0	19.50
	B	1.8	1.10
	C	0.1	0.30
Round 2	A	49	0.10
	B	5.2	0.20
	C	0.3	0.04

Phagemid DNA from 12 colonies from the second round of plating together with 5 colonies from a plating of the ini-

tial phage mixtures was isolated and the nucleotide sequence of the CTLD region determined. From the initial 1/10 mixture (B series) of PhtCTLD/PhtCPB one out of five were identified as the CTLD sequence. From the initial 1/10⁵ mixture (C series) all five sequences were derived from the pPhtCPB phagemid. After round 2 nine of the twelve sequences analysed from the B series and all twelve sequences from the C series were derived from the pPhtCTLD phagemid.

10 Example 8

Model selection of CTLD-phages on plasminogen.

II: elution by 0.1 M triethylamine after panning.

In order to demonstrate that tetranectin derived CTLD-bearing phages can be selected from a population of phages, mixtures of PhtCTLD phages isolated from a *E. coli* TG1 culture transformed with the phagemid pPhtCTLD (cf, EXAMPLE 1) after infection with M13K07 helper phage and phages isolated from a culture transformed with the phagemid pPhtCPB (cf, EXAMPLE 6) after infection with M13K07 helper phage at ratios of 1:10² and 1:10⁶, respectively were used in a selection experiment using panning in 96-well Maxisorb microtiterplates (NUNC, Denmark) and with human plasminogen as antigen using standard procedures.

Briefly, in each well 3 µg of human plasminogen in 100 µL PBS (PBS, 0.2 g KCl, 0.2 g KH₂PO₄, 8 g NaCl, 1.44 g Na₂HPO₄, 2H₂O, water to 1 L, and adjusted to pH 7.4 with NaOH) or 100 µL PBS (for analysis of non specific binding) was used for over night coating at 4 °C and at 37 °C for one hour. After washing once with PBS, wells were blocked with 400 µL PBS and 3% non fat dried milk for one hour at 37 °C. After blocking wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of phages suspended in 100 µL PBS, 3% non fat dried milk. The phages

were allowed to bind at 37 °C for one hour before washing 15 times with PBS, Tween 20, and 15 times with PBS. Bound phages were eluted from each well by 100 µL 0.1 M triethylamine for 10 min at room temperature, and upon neutralisation with 0.5 vol. 1 M Tris-HCl pH 7.4, used for infection of exponentially growing *E. coli* TG1 cells before plating and titration on 2xTY agar plates containing 2% glucose and 0.1 mg/mL ampicillin.

Initially (round 1) 10^{12} PhtCTLD phages (A series), a mixture of 10^9 PhtCTLD phages and 10^{11} PhtCPB phages (B series), or a mixture of 10^5 PhtCTLD and 10^{11} PhtCPB phages (C series) were used. In the following round (round 2) 10^{11} phages of the output from each series were used. Results from the two rounds of selection are summarised in Table 5.

Table 5: Selection of mixtures of PhtCTLD and PhtCPB by panning elution with triethylamine.

		Plasminogen (* 10^4 colonies)	Blank (* 10^4 colonies)
Round 1	A	18	0.02
	B	0.5	0.00
	C	0.25	0.02
Round 2	A	n.d.	n.d.
	B	5.0	0.00
	C	1.8	0.02
Round 3	A	n.d.	n.d.
	B	11	0.00
	C	6.5	0.02

n.d. = not determined

Phage mixtures from the A and the B series from the second round of selection were grown using a standard procedure, and analysed for binding to plasminogen in an ELISA-type assay. Briefly, in each well 3 μ g of plasminogen in 100 μ L
5 PBS (PBS, 0.2 g KCl, 0.2 g KH₂PO₄, 8 g NaCl, 1.44 g Na₂HPO₄, 2H₂O, water to 1 L, and adjusted to pH 7.4 with NaOH) or 100 μ L PBS (for analysis of non specific binding) was used for over night coating at 4 °C and at 37 °C for one hour. After washing once with PBS, wells were blocked
10 with 400 μ L PBS and 3% non fat dried milk for one hour at 37°C. After blocking wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of phages suspended in 100 μ L PBS, 3% non fat dried milk. The phage mixtures were allowed to bind at 37 °C for one hour
15 before washing three times with PBS, Tween 20, and three times with PBS. After washing, 50 μ L of a 1:5000 dilution of a HRP-conjugated anti-gene VIII antibody (Amersham Pharmacia Biotech) in PBS, 3% non fat dried milk was added to each well and incubated at 37 °C for one hour. After bind-
20 ing of the "secondary" antibody wells were washed three times with PBS, Tween 20, and three times with PBS before the addition of 50 μ L of TMB substrate (DAKO-TMB One-Step Substrate System, code: S1600, DAKO, Denmark). Reaction was allowed to proceed for 20 min. before quenching with 0.5
25 vol. 0.5 M H₂SO₄, and analysis. The result of the ELISA analysis confirmed specific binding to plasminogen of phages in both series (fig. 28).

Example 9

30 Selection of phages from the library Phtlec-lb002 binding to hen egg white lysozyme.

1.2*10¹² phages, approximately 250 times the size of the original library, derived from the Phtlec-lb002 library

(cf, EXAMPLE 4) were used in an experimental procedure for the selection of phages binding to hen egg white lysozyme involving sequential rounds of panning using standard procedures.

5 Briefly, 30 μ g of hen egg white lysozyme in 1 mL PBS (PBS, 0.2 g KCl, 0.2 g KH_2PO_4 , 8 g NaCl, 1.44 g Na_2HPO_4 , $2\text{H}_2\text{O}$, water to 1 L, and adjusted to pH 7.4 with NaOH) or 1 mL PBS (for analysis of non specific binding) was used for over
10 night coating of Maxisorb immunotubes (NUNC, Denmark) at 4 $^{\circ}\text{C}$ and at 37 $^{\circ}\text{C}$ for one hour. After washing once with PBS, tubes were filled and blocked with PBS and 3% non fat dried milk for one hour at 37 $^{\circ}\text{C}$. After blocking tubes were washed once in PBS, 0.1% Tween 20 and three times with PBS before the addition of phages suspended in 1 mL PBS, 3% non fat
15 dried milk. The phages were allowed to bind at 37 $^{\circ}\text{C}$ for one hour before washing six times with PBS, Tween 20 and six times with PBS. Bound phages were eluted from each well by 1 mL 0.1 M triethylamine for 10 min at room temperature, and upon neutralisation with 1 M Tris-HCl pH 7.4, used for
20 infection of exponentially growing *E. coli* TG1 cells before plating and titration on 2xTY agar plates containing 2% glucose and 0.1 mg/mL ampicillin. In the subsequent rounds of selection approximately 10^{12} phages derived from a culture grown from the colonies plated after infection with
25 the phages eluted from the lysozyme coated tube were used in the panning procedure. However, the stringency in binding was increased by increasing the number of washing step after phage panning from six to ten.

The results from the selection procedure is shown in Table
30 7.

Table 7: Selection by panning of lysozyme binding phages from Phtlec-lb002 library.

	Lysozyme	Blank	Ratio
Round 1	$2.4 \cdot 10^4$	n.a.	n.a.
Round 2	$3.5 \cdot 10^3$	$4.0 \cdot 10^2$	9
Round 3	$3.2 \cdot 10^5$	$2.5 \cdot 10^2$	$1.3 \cdot 10^3$

n.a. = not applicable

Phages were grown from twelve clones isolated from the third round of selection in order to analyse the specificity of binding using a standard procedure, and analysed for binding to hen egg white lysozyme and human β_2 -microglobulin in an ELISA-type assay. Briefly, in each well 3 μ g of hen egg white lysozyme in 100 μ L PBS (PBS, 0.2 g KCl, 0.2 g KH_2PO_4 , 8 g NaCl, 1.44 g Na_2HPO_4 , $2\text{H}_2\text{O}$, water to 1 L, and adjusted to pH 7.4 with NaOH), or 3 μ g of human β_2 -microglobulin, or 100 μ L PBS (for analysis of non specific binding) was used for over night coating at 4 $^\circ\text{C}$ and at 37 $^\circ\text{C}$ for one hour. After washing once with PBS, wells were blocked with 400 μ L PBS and 3% non fat dried milk for one hour at 37 $^\circ\text{C}$. After blocking wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of phages suspended in 100 μ L PBS, 3% non fat dried milk. The phages were allowed to bind at 37 $^\circ\text{C}$ for one hour before washing three times with PBS, Tween 20 and three times with PBS. After washing, 50 μ L of a 1 to 5000 dilution of a HRP-conjugated anti-gene VIII antibody (Amersham Pharmacia Biotech) in PBS, 3% non fat dried milk was added to each well and incubated at 37 $^\circ\text{C}$ for one hour. After binding of the "secondary" antibody wells were washed three times with PBS, Tween 20 and three times with PBS before the addition of 50 μ L of TMB substrate (DAKO-TMB One-Step Substrate System, code: S1600, DAKO, Denmark): Reac-

tion was allowed to proceed for 20 min before quenching with 0.5 M H₂SO₄.

Results showing relatively weak but specific binding to lysozyme are summarised in Fig. 29.

5

EXAMPLE 10

Construction of the rat mannose-binding protein CTLD (r-MBP) derived phagemid (pPrMBP) and human lung surfactant protein D CTLD (h-SP-D) derived phagemid (pPhSP-D)

- 10 The phagemid, pPrMBP, is constructed by ligation of the *Sfi* I and Not I restricted DNA fragment amplified from cDNA, isolated from rat liver (Drickamer, K., et al., *J. Biol. Chem.* 1987, 262(6):2582-2589) (with the oligonucleotide primers SfiMBP 5'-CGGCTGAGCGGCCCGCCGATGGC-
- 15 CGAGCCAAACAAGTTGCATGCCTTCTCC-3' [SEQ ID NO:62] and NotMBP 5'-GCACTCCTGCGGCCGCGCTGGGAACTCGCAGAC-3' [SEQ ID NO:63]) into a *Sfi* I and Not I precut vector, pCANTAB 5E supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) using standard procedures. Outlines of the resulting pPrMBP is
- 20 shown in Fig. 31 and the nucleotide sequence of PrMBP is given as (SEQ ID NO:58). The amino acid sequence encoded by the PrMBP insert is shown in Fig. 30 (SEQ ID NO:59).

- The phagemid, pPhSP-D, is constructed by ligation of the *Sfi* I and Not I restricted DNA fragment amplified from cDNA,
- 25 isolated from human lung (Lu, J., et al., *Biochem J.* 1992 Jun 15; 284:795-802) (with the oligonucleotide primers SfiSP-D 5'-CGGCTGAGCGGCCCGCCGATGGCCGAGCCAAAGAAAGTTGAGCTCTTCCC-3' [SEQ ID NO:64] and NotSP-D 5'-GCACTCCTGCGGCCGCGAACTCGCAGACCACAAGAC-3' [SEQ ID NO:65]) into a *Sfi* I and
- 30 Not I precut vector, pCANTAB 5E supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) using standard procedures. Outlines of the resulting pPhSP-D is shown in Fig.

33 and the nucleotide sequence of PhSP-D, is given as (SEQ ID NO:60). The amino acid sequences encoded by the PhSP-D insert is shown in Fig 32 (SEQ ID NO:61).

5 Example 11

Construction of the phage library PrMBP-lb001

The phage library PrMBP-lb001, containing random amino acid residues corresponding to PrMBP CTLD (SEQ ID NO:59) positions 71 to 73 or 70 to 76 (loop 1) and 97 to 101 or 100 to 101 (loop 4) is constructed by ligation of 20 µg *Sfi*I and NotI restricted pPrMBP phagemid DNA (cf. Example 10) with 10 µg of a *Sfi*I and NotI restricted DNA fragment population encoding the appropriately randomised loop 1 and 4 regions. The DNA fragment population is amplified, from nine assembly reactions combining each of the three loop 1 DNA fragments with each of the three loop 4 DNA fragments as templates and the oligonucleotides *Sfi*-tag 5'-CGGCTGAGCGGCCCA-GC-3' (SEQ ID NO:74) and Not-tag 5'-GCACTCCTGCGGCCGCG-3' (SEQ ID NO:75) as primers using standard procedures.

Each of the three loop 1 fragments is amplified in a primary PCR reaction with pPrMBP phagemid DNA (cf. Example 10) as template and the oligonucleotides MBPloop1a fo (SEQ ID NO:66), MBPloop1b fo (SEQ ID NO:67) or MBPloop1c fo (SEQ ID NO:68) and *Sfi*MBP (SEQ ID NO:62) as primers, and further amplified in a secondary PCR reaction using *Sfi*-tag (SEQ ID NO:74) and MBPloop1-tag fo (SEQ ID NO:69). Each of the three DNA loop 4 fragments is amplified in a primary PCR reaction with pPrMBP phagemid DNA (cf. Example 10) as template and the oligonucleotides MBPloop4a rev (SEQ ID NO:71), MBPloop4b rev (SEQ ID NO:72) or MBPloop4c rev (SEQ ID NO:73) and NotMBP (SEQ ID NO:63) as primers using standard procedures and further amplified in a secondary PCR reaction using MBPloop4-tag rev (SEQ ID NO:70) and Not-tag

(SEQ ID NO:63). In the oligonucleotide sequences N denotes a mixture of 25% of each of the nucleotides T, C, G, and A, respectively, and S denotes a mixture of 50 % of C and G, encoding the appropriately randomized nucleotide sequence.

5 The ligation mixture is used to transform so-called electrocompetent *E. coli* TG-1 cells by electroporation using standard procedures. After transformation the *E. coli* TG-1 cells are plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated over night at 30

10 °C.

Example 12

Construction of the phage library PhSP-D-1b001

The phage library PhSP-D-1b001, containing random amino

15 acid residues corresponding to PhSP-D CTLD insert (SEQ ID NO:61) positions 74 to 76 or 73 to 79 (loop 1) and 100 to 104 or 103 to 104 (loop 4) is constructed by ligation of 20 µg *Sfi*I and *Not*I restricted pPhSP-D phagemid DNA (cf. Example 10) with 10 µg of a *Sfi*I and *Not*I restricted DNA fragment

20 population encoding the appropriately randomised loop 1 and 4 regions. The DNA fragment population is amplified, from nine assembly reactions combining each of the three loop 1 DNA fragments with each of the three loop 4 DNA fragments as templates and the oligonucleotides *Sfi*-tag 5'-

25 CGGCTGAGCGGCCAGC-3' (SEQ ID NO:74) and *Not*-tag 5'- GCACTCCTGCGGCCGCG-3' (SEQ ID NO:75) as primers using standard procedures. Each of the three loop 1 fragments is amplified in a primary PCR reaction with pPhSP-D phagemid DNA (cf. Example 10) as template and the oligonucleotides *Sp*-

30 *dloop1a* fo (SEQ ID NO:76), *Sp-dloop1b* fo (SEQ ID NO:77) or *Sp-dloop1c* fo (SEQ ID NO:78) and *Sfi*SP-D (SEQ ID NO:64) as primers, and further amplified in a PCR reaction using *Sfi*-tag (SEQ ID NO:74) and *Sp-dloop1-tag* fo (SEQ ID NO:79) as

primers. Each of the three DNA loop 4 fragments is amplified in a primary PCR reaction with pPhSP-D phagemid DNA (cf. Example 10) as template and the oligonucleotides Sp-dloop4a rev (SEQ ID NO:81), Sp-dloop4b rev (SEQ ID NO:82) or Sp-dloop4c rev (SEQ ID NO:83) and NotSP-D (SEQ ID NO:65) as primers using standard procedures and further amplified in a PCR reaction using Sp-dloop4-tag rev (SEQ ID NO:80) and Not-tag (SEQ ID NO:75) as primers. In the oligonucleotide sequences N denotes a mixture of 25% of each of the nucleotides T, C, G, and A, respectively, and S denotes a mixture of 50 % of C and G, encoding the appropriately randomized nucleotide sequence. The ligation mixture is used to transform so-called electrocompetent *E. coli* TG-1 cells by electroporation using standard procedures. After transformation the *E. coli* TG-1 cells are plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated over night at 30 °C.

Example 13

20 Construction of the phage library PhtCTLD-lb004

All oligonucleotides used in this example were supplied by DNA Technology (Aarhus, Denmark).

The phage library PhtCTLD-lb004, containing random amino acid residues corresponding to PhtCTLD (SEQ ID NO:15) positions 97 to 102 or 98 to 101(loop 3) and positions 116 to 122 or 118 to 120 (loop 5) was constructed by ligation of 20 µg *Kpn*I and *Mun*I restricted pPhtCTLD phagemid DNA (cf. Example 1) with 10 µg of a *Kpn*I and *Mun*I restricted DNA fragment population encoding the randomised loop 3 and 5 regions. The DNA fragment population was amplified from nine primary PCR reactions combining each of the three loop 3 DNA fragments with each of the three loop 5 DNA fragments. The fragments was amplified with either of the oli-

gonucleotides loop3a (SEQ ID NO:84), loop3b (SEQ ID NO: 85), or loop3c (SEQ ID NO:86) as template and loop5a(SEQ ID NO:87), loop5b(SEQ ID NO:88)or loop5c(SEQ ID NO:89) and loop3-4rev(SEQ ID NO:91) as primers. The DNA fragments were
5 further amplified in PCR reactions, using the primary PCR product as template and the oligonucleotide loop3-4rev (SEQ ID NO:91) and loop3-4-5tag fo (SEQ ID NO:90) as primers. All PCR reactions were performed using standard procedures.

In the oligonucleotide sequences N denotes a mixture of 25%
10 of each of the nucleotides T, C, G, and A, respectively and S denotes a mixture of 50 % of C and G, encoding the appropriately randomised nucleotide sequence. The ligation mixture was used to transform so-called electrocompetent E. coli TG-1 cells by electroporation using standard proce-
15 dures. After transformation the E. coli TG-1 cells were plated on 2xTY-agar plates containing 0.2 mg ampicillin/mL and 2% glucose and incubated over night at 30 °C.

The size of the resulting library, PhtCTLD-lb004, was determined to $7 \cdot 10^9$ clones. Sixteen clones from the library
20 were picked and phagemid DNA isolated. The nucleotide sequence of the loop-regions were determined (DNA Technology, Aarhus, Denmark). Thirteen clones were found to contain correct loop inserts and three clones contained a frameshift mutation in the region.

25

Example 14

Selection of Phtlec-phages and PhtCTLD-phages binding to the blood group A sugar moiety immobilised on human serum albumin

30 Phages grown from glycerol stocks of the libraries Phtlec-lb001 and Phtlec-lb002 (cf. Example 4) and phages grown from a glycerol stock of the library PhtCTLD-lb003 (cf. Ex-

ample 5), using a standard procedure, were used in an experiment designed for the selection of Phtlec- and PhtCTLD derived phages with specific affinity to the blood group A sugar moiety immobilized on human serum albumin, A-HA, by
5 panning in 96-well Maxisorb micro-titerplates (NUNC, Denmark) using standard procedures.

Initially, the phage supernatants were precipitated with 0.3 vol. of a solution of 20% polyethylene glycol 6000 (PEG) and 2.5 M NaCl, and the pellets re-suspended in TE-
10 buffer (10 mM Tris-HCl pH 8, 1 mM EDTA). After titration on *E. coli* TG-1 cells, phages derived from Phtlec-lb001 and -lb002 were mixed (#1) in a 1:1 ratio and adjusted to 5×10^{12} pfu/mL in 2*TY medium, and phages grown from the PhtCTLD-lb003 library (#4) were adjusted to 2.5×10^{12} pfu/mL in 2*TY
15 medium.

One microgram of the "antigen", human blood group A trisaccharide immobilised on human serum albumin, A-HA, (Glycorex AB, Lund, Sweden) in 100 μ L PBS (PBS, 0.2 g KCl, 0.2 g KH_2PO_4 , 8 g NaCl, 1.44 g $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, water to 1 L, and
20 adjusted to pH 7.4 with NaOH), in each of three wells, was coated over night at 4 °C and at room temperature for one hour, before the first round of panning. After washing once with PBS, wells were blocked with 300 μ L PBS and 3% non fat dried milk for one hour at room temperature. After blocking
25 wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of a mixture of 50 μ L of the phage suspension and 50 μ L PBS, 6% non fat dried milk. The phages were allowed to bind at room temperature for two hours before washing eight times with PBS, Tween 20, and
30 eight times with PBS. Bound phages were eluted from each well by trypsin digestion in 100 μ L (1 mg/mL trypsin in PBS) for 30 min. at room temperature, and used for infection of exponentially growing *E. coli* TG1 cells before

plating and titration on 2xTY agar plates containing 2% glucose and 0.1 mg/mL ampicillin.

In the second round of selection, 150 μ L of crude phage supernatant, grown from the first round output colonies, was
5 mixed with 150 μ L PBS, 6% non fat dried milk, and used for panning distributing 100 μ L of the mixture in each of three A-HA coated wells, as previously described. Stringency in binding was increased by increasing the number of washing steps from 16 to 32. 300 μ L of phage mixture was also used
10 for panning in three wells, which had received no antigen as control.

In the third round of selection, 150 μ L of crude phage supernatant, grown from the second round output colonies, was mixed with 150 μ L PBS, 6% non fat dried milk, and used for
15 panning distributing 100 μ L of the mixture in each of three A-HA coated wells, as previously described. The number of washing steps was again 32. 300 μ L of phage mixture was also used for panning in three wells, which had received no antigen as control.

20 The results from the selection procedure are summarised in Table 8

Table 8. Selection of Phtlec phages (#1) and PhtCTLD phages (#4) binding to A-HA by panning and elution with trypsin digestion.

		A-HA	Blank	Ratio
Round 1	#1	0.8×10^3	n.a.	n.a.
	#4	1.1×10^3	n.a.	n.a.
Round 2	#1	1.0×10^3	0.5×10^2	20
	#4	1.3×10^3	0.5×10^2	26
Round 3	#1	8.0×10^4	0.5×10^2	1600
	#4	9.0×10^5	0.5×10^2	18000

n.a. not applicable.

- 5 48 clones from each of the #1 and #4 series were picked and grown in a 96 well microtiter tray and phages produced by infection with M13K07 helper phage using a standard procedure. Phages from the 96 phage supernatants were analysed for binding to the A-HA antigen and for non-specific binding to hen egg white lysozyme using an ELISA-type assay.
- 10 Briefly, in each well 1 μ g of A-HA in 100 μ L PBS (PBS, 0.2 g KCl, 0.2 g KH_2PO_4 , 8 g NaCl, 1.44 g Na_2HPO_4 , $2\text{H}_2\text{O}$, water to 1 L, and adjusted to pH 7.4 with NaOH) or 1 μ g of hen egg white lysozyme in 100 μ L PBS (for analysis of non specific binding) was used for over night coating at 4 $^\circ\text{C}$ and
- 15 at room temperature for one hour. After washing once with PBS, wells were blocked with 300 μ L PBS and 3% non fat dried milk for one hour at room temperature. After blocking wells were washed once in PBS and 0.1% Tween 20 and three times with PBS before the addition of 50 μ L phage supernatant in 50 μ L PBS, 6% non fat dried milk. The phage mixtures were allowed to bind at room temperature for two hours before washing three times with PBS, Tween 20, and three times with PBS. After washing, 50 μ L of a 1:5000 di-
- 20

lution of a HRP-conjugated anti-gene VIII antibody (Amersham Pharmacia Biotech) in PBS, 3% non fat dried milk, was added to each well and incubated at room temperature for one hour. After binding of the "secondary" antibody wells
5 were washed three times with PBS, Tween 20, and three times with PBS before the addition of 50 μ L of TMB substrate (DAKO-TMB One-Step Substrate System, DAKO, Denmark). Reaction was allowed to proceed for 20 min. before quenching with 0.5 M H_2SO_4 , and analysis. The result of the ELISA
10 analysis showed "hits" in terms of specific binding to A-HA of phages in both series (fig. 34 and 35), as judged by a signal ratio between signal on A-HA to signal on lysozyme at or above 1.5, and with a signal above background.

From the #1 series 13 hits were identified and 28 hits were
15 identified from the #4 series.

REFERENCES

- Aspberg, A., Miura, R., Bourdoulous, S., Shimonaka, M., Heinegård, D., Schachner, M., Ruoslahti, E., and Yamaguchi, Y. (1997). "The C-type lectin domains of lecticans, a family of aggregating chondroitin sulfate proteoglycans, bind
20 tenascin-R by protein-protein interactions independent of carbohydrate moiety". *Proc. Natl. Acad. Sci. (USA)* 94: 10116-10121
- Bass, S., Greene, R., and Wells, J.A. (1990). "Hormone
25 phage: an enrichment method for variant proteins with altered binding properties". *Proteins* 8: 309-314
- Benhar, I., Azriel, R., Nahary, L., Shaky, S., Berdichevsky, Y., Tamarkin, A., and Wels, W. (2000). "Highly efficient selection of phage antibodies mediated by display of
30 antigen as Lpp-OmpA' fusions on live bacteria". *J. Mol. Biol.* 301: 893-904

- Berglund, L. and Petersen, T.E. (1992). "The gene structure of tetranectin, a plasminogen binding protein". *FEBS Letters* 309: 15-19
- 5 Bertrand, J.A., Pignol, D., Bernard, J-P., Verdier, J-M., Dagorn, J-C., and Fontecilla-Camps, J.C. (1996). "Crystal structure of human lithostathine, the pancreatic inhibitor of stone formation". *EMBO J.* 15: 2678-2684
- 10 Bettler, B., Texido, G., Raggini, S., Rüegg, D., and Hofstetter, H. (1992). "Immunoglobulin E-binding site in Fc epsilon receptor (Fc epsilon RII/CD23) identified by homolog-scanning mutagenesis". *J. Biol. Chem.* 267: 185-191
- 15 Blanck, O., Iobst, S.T., Gabel, C., and Drickamer, K. (1996). "Introduction of selectin-like binding specificity into a homologous mannose-binding protein". *J. Biol. Chem.* 271: 7289-7292
- Boder, E.T. and Wittrup, K.D. (1997). "Yeast surface display for screening combinatorial polypeptide libraries". *Nature Biotech.* 15: 553-557
- 20 Burrows L, Iobst ST, Drickamer K. (1997) "Selective binding of N-acetylglucosamine to the chicken hepatic lectin". *Biochem J.* 324:673-680
- 25 Chiba, H., Sano, H., Saitoh, M., Sohma, H., Voelker, D.R., Akino, T., and Kuroki, Y. (1999). "Introduction of mannose binding protein-type phosphatidylinositol recognition into pulmonary surfactant protein A". *Biochemistry* 38: 7321-7331
- 30 Christensen, J.H., Hansen, P.K., Lillelund, O., and Thøgersen, H.C. (1991). "Sequence-specific binding of the N-terminal three-finger fragment of *Xenopus* transcription factor IIIA to the internal control region of a 5S RNA gene". *FEBS Letters* 281: 181-184

- Cyr, J.L. and Hudspeth, A.J. (2000). "A library of bacteriophage-displayed antibody fragments directed against proteins of the inner ear". *Proc. Natl. Acad. Sci (USA)* 97: 2276-2281
- 5 Drickamer, K. (1992). "Engineering galactose-binding activity into a C-type mannose-binding protein". *Nature* 360: 183-186
- Drickamer, K. and Taylor, M.E. (1993). "Biology of animal lectins". *Annu. Rev. Cell Biol.* 9: 237-264
- 10 Drickamer, K. (1999). "C-type lectin-like domains". *Curr. Opinion Struc. Biol.* 9: 585-590
- Dunn, I.S. (1996). "Phage display of proteins". *Curr. Opinion Biotech.* 7: 547-553
- 15 Erbe, D.V., Lasky, L.A., and Presta, L.G. "Selectin variants". *US Patent No. 5593882*
- Ernst, W.J., Spenger, A., Toellner, L., Katinger, H., Grabherr, R.M. (2000). "Expanding baculovirus surface display. Modification of the native coat protein gp64 of *Autographa californica* NPV". *Eur. J. Biochem.* 267: 4033-
20 4039
- Ewart, K.V., Li, Z., Yang, D.S.C., Fletcher, G.L., and Hew, C.L. (1998). "The ice-binding site of Atlantic herring antifreeze protein corresponds to the carbohydrate-binding site of C-type lectins". *Biochemistry* 37: 4080-4085
- 25 Feinberg, H., Park-Snyder, S., Kolatkar, A.R., Heise, C.T., Taylor, M.E., and Weis, W.I. (2000). "Structure of a C-type carbohydrate recognition domain from the macrophage mannose receptor". *J. Biol. Chem.* 275: 21539-21548

- Fujii, I., Fukuyama, S., Iwabuchi, Y., and Tanimura, R. (1998). "Evolving catalytic antibodies in a phage-displayed combinatorial library". *Nature Biotech.* 16: 463-467
- Gates, C.M., Stemmer, W.P.C., Kaptein, R., and Schatz, P.J. 5 (1996). "Affinity selective isolation of ligands from peptide libraries through display on a lac repressor "head-piece dimer". *J. Mol. Biol.* 255: 373-386
- Graversen, J.H., Lorentsen, R.H., Jacobsen, C., Moestrup, S.K., Sigurskjold, B.W., Thøgersen, H.C., and Etzerodt, M. 10 (1998). "The plasminogen binding site of the C-type lectin tetranectin is located in the carbohydrate recognition domain, and binding is sensitive to both calcium and lysine". *J. Biol. Chem.* 273:29241-29246
- Graversen, J.H., Jacobsen, C., Sigurskjold, B.W., Lorentsen, R.H., Moestrup, S.K., Thøgersen, H.C., and Etzerodt, M. 15 (2000). "Mutational Analysis of Affinity and Selectivity of Kringle-Tetranectin Interaction. Grafting novel kringle affinity onto the tetranectin lectin scaffold". *J. Biol. Chem.* 275: 37390-37396
- Griffiths, A.D. and Duncan, A.R. 20 (1998). "Strategies for selection of antibodies by phage display". *Curr. Opinion Biotech.* 9: 102-108
- Holtet, T.L., Graversen, J.H., Clemmensen, I., Thøgersen, H.C., and Etzerodt, M. 25 (1997). "Tetranectin, a trimeric plasminogen-binding C-type lectin". *Prot. Sci.* 6: 1511-1515
- Honma, T., Kuroki, Y., Tzunezawa, W., Ogasawara, Y., Sohma, H., Voelker, D.R., and Akino, T. (1997). "The mannose-binding protein A region of glutamic acid185-alanine221 can functionally replace the surfactant protein A region of 30 glutamic acid195-phenylalanine228 without loss of interac-

- tion with lipids and alveolar type II cells". *Biochemistry* 36: 7176-7184
- Huang, W., Zhang, Z., and Palzkill, T. (2000). "Design of potent beta-lactamase inhibitors by phage display of beta-lactamase inhibitory protein". *J. Biol. Chem.* 275: 14964-14968
- Hufton, S.E., van Neer, N., van den Beuken, T., Desmet, J., Sablon, E., and Hoogenboom, H.R. (2000). "Development and application of cytotoxic T lymphocyte-associated antigen 4 as a protein scaffold for the generation of novel binding ligands". *FEBS Letters* 475: 225-231
- Håkansson, K., Lim, N.K., Hoppe, H-J., and Reid, K.B.M. (1999). "Crystal structure of the trimeric alpha-helical coiled-coil and the three lectin domains of human lung surfactant protein D". *Structure Folding and Design* 7: 255-264
- Iobst, S.T., Wormald, M.R., Weis, W.I., Dwek, R.A., and Drickamer, K. (1994). "Binding of sugar ligands to Ca(2+)-dependent animal lectins. I. Analysis of mannose binding by site-directed mutagenesis and NMR". *J. Biol. Chem.* 269: 15505-15511
- Iobst, S.T. and Drickamer, K. (1994). "Binding of sugar ligands to Ca(2+)-dependent animal lectins. II. Generation of high-affinity galactose binding by site-directed mutagenesis". *J. Biol. Chem.* 269: 15512-15519
- Iobst, S.T. and Drickamer, K. (1996). "Selective sugar binding to the carbohydrate recognition domains of the rat hepatic and macrophage asialoglycoprotein receptors". *J. Biol. Chem.* 271: 6686-6693
- Jaquinod, M., Holtet, T. L., Etzerodt, M., Clemmensen, I., Thøgersen, H. C., and Roepstorff, P. (1999). "Mass Spectro-

- metric Characterisation of Post-Translational Modification and Genetic Variation in Human Tetranectin". *Biol. Chem.* 380: 1307-1314
- 5 Kastrup, J.S., Nielsen, B.B., Rasmussen, H., Holtet, T.L., Graversen, J.H., Etzerodt, M., Thøgersen, H.C., and Larsen, I.K. (1998). "Structure of the C-type lectin carbohydrate recognition domain of human tetranectin". *Acta. Cryst. D* 54: 757-766
- 10 Kogan, T.P., Revelle, B.M., Tapp, S., Scott, D., and Beck, P.J. (1995). "A single amino acid residue can determine the ligand specificity of E-selectin". *J. Biol. Chem.* 270: 14047-14055
- 15 Kolatkar, A.R., Leung, A.K., Isecke, R., Brossmer, R., Drickamer, K., and Weis, W.I. (1998). "Mechanism of N-acetylgalactosamine binding to a C-type animal lectin carbohydrate-recognition domain". *J. Biol. Chem.* 273: 19502-19508
- 20 Lorentsen, R.H., Graversen, J.H., Caterer, N.R., Thøgersen, H.C., and Etzerodt, M. (2000). "The heparin-binding site in tetranectin is located in the N-terminal region and binding does not involve the carbohydrate recognition domain". *Biochem. J.* 347: 83-87
- 25 Marks, J.D., Hoogenboom, H.R., Griffiths, A.D., and Winter, G. (1992). "Molecular evolution of proteins on filamentous phage. Mimicking the strategy of the immune system". *J. Biol. Chem.* 267: 16007-16010
- 30 Mann K, Weiss IM, Andre S, Gabius HJ, Fritz M. (2000). "The amino-acid sequence of the abalone (*Haliotis laevis*) na-cre protein perlucin. Detection of a functional C-type lectin domain with galactose/mannose specificity". *Eur. J. Biochem.* 267: 5257-5264

- McCafferty, J., Jackson, R.H., and Chiswell, D.J. (1991). "Phage-enzymes: expression and affinity chromatography of functional alkaline phosphatase on the surface of bacteriophage". *Prot. Eng.* 4: 955-961
- 5 McCormack, F.X., Kuroki, Y., Stewart, J.J., Mason, R.J., and Voelker, D.R. (1994). "Surfactant protein A amino acids Glu195 and Arg197 are essential for receptor binding, phospholipid aggregation, regulation of secretion, and the facilitated uptake of phospholipid by type II cells". *J. Biol. Chem.* 269: 29801-29807
- 10 McCormack, F.X., Festa, A.L., Andrews, R.P., Linke, M., and Walzer, P.D. (1997). "The carbohydrate recognition domain of surfactant protein A mediates binding to the major surface glycoprotein of *Pneumocystis carinii*". *Biochemistry* 36: 8092-8099
- 15 Meier, M., Bider, M.D., Malashkevich, V.N., Spiess, M., and Burkhard, P. (2000). "Crystal structure of the carbohydrate recognition domain of the H1 subunit of the asialoglycoprotein receptor". *J. Mol. Biol.* 300: 857-865
- 20 Mikawa, Y.G., Maruyama, I.N., and Brenner, S. (1996). "Surface display of proteins on bacteriophage lambda heads". *J. Mol. Biol.* 262: 21-30
- Mio H, Kagami N, Yokokawa S, Kawai H, Nakagawa S, Takeuchi K, Sekine S, Hiraoka A. (1998). "Isolation and characterization of a cDNA for human mouse, and rat full-length stem cell growth factor, a new member of C-type lectin superfamily". *Biochem. Biophys. Res. Commun.* 249: 124-130
- 25 Mizuno, H., Fujimoto, Z., Koizumi, M., Kano, H., Atoda, H., and Morita, T. (1997). "Structure of coagulation factors IX/X-binding protein, a heterodimer of C-type lectin domains". *Nat. Struc. Biol.* 4: 438-441
- 30

- Ng, K.K., Park-Snyder, S., and Weis, W.I. (1998a). "Ca²⁺-dependent structural changes in C-type mannose-binding proteins". *Biochemistry* 37: 17965-17976
- Ng, K.K. and Weis, W.I. (1998b). "Coupling of prolyl peptide bond isomerization and Ca²⁺ binding in a C-type mannose-binding protein". *Biochemistry* 37: 17977-17989
- Nielsen, B.B., Kastrup, J.S., Rasmussen, H., Holtet, T.L., Graversen, J.H., Etzerodt, M., Thøgersen, H.C., and Larsen, I.K. (1997). "Crystal structure of tetranectin, a trimeric plasminogen-binding protein with an alpha-helical coiled coil". *FEBS Letters* 412: 388-396
- Nissim A., Hoogenboom, H.R., Tomlinson, I.M., Flynn, G., Midgley, C., Lane, D., and Winter, G. (1994). "Antibody fragments from a 'single pot' phage display library as immunochemical reagents". *EMBO J.* 13: 692-698
- Ogasawara, Y. and Voelker, D.R. (1995). "Altered carbohydrate recognition specificity engineered into surfactant protein D reveals different binding mechanisms for phosphatidylinositol and glucosylceramide". *J. Biol. Chem.* 270: 14725-14732
- Ohtani, K., Suzuki, Y., Eda, S., Takao, K., Kase, T., Yamazaki, H., Shimada, T., Keshi, H., Sakai, Y., Fukuoh, A., Sakamoto, T., and Wakamiya, N. (1999). "Molecular cloning of a novel human collectin from liver (CL-L1)". *J. Biol. Chem.* 274: 13681-13689
- Pattanajitvilai, S., Kuroki, Y., Tsunazawa, W., McCormack, F.X., and Voelker, D.R. (1998). "Mutational analysis of Arg197 of rat surfactant protein A. His197 creates specific lipid uptake defects". *J. Biol. Chem.* 273: 5702-5707

- Poget, S.F., Legge, G.B., Proctor, M.R., Butler, P.J., Bycroft, M., and Williams, R.L. (1999). "The structure of a tunicate C-type lectin from *Polyandrocarpa misakiensis* complexed with D-galactose". *J. Mol. Biol.* 290: 867-879
- 5 Revelle, B.M., Scott, D., Kogan, T.P., Zheng, J., and Beck, P.J. (1996). "Structure-function analysis of P-selectin-sialyl LewisX binding interactions. Mutagenic alteration of ligand binding specificity". *J. Biol. Chem.* 271: 4289-4297
- 10 Sano, H., Kuroki, Y., Honma, T., Ogasawara, Y., Sohma, H., Voelker, D.R., and Akino, T. (1998). "Analysis of chimeric proteins identifies the regions in the carbohydrate recognition domains of rat lung collectins that are essential for interactions with phospholipids, glycolipids, and alveolar type II cells". *J. Biol. Chem.* 273: 4783-4789
- 15 Schaffitzel, C., Hanes, J., Jermutus, L., and Plückert, A. (1999). "Ribosome display: an in vitro method for selection and evolution of antibodies from libraries". *J. Immunol. Methods* 231: 119-135
- 20 Sheriff, S., Chang, C.Y., and Ezekowitz, R.A. (1994). "Human mannose-binding protein carbohydrate recognition domain trimerizes through a triple alpha-helical coiled-coil". *Nat. Struct. Biol.* 1: 789-794
- 25 Sørensen, C.B., Berglund, L., and Petersen, T.E. (1995). "Cloning of a cDNA encoding murine tetranectin". *Gene* 152: 243-245
- Torgersen, D., Mullin, N.P., and Drickamer, K. (1998). "Mechanism of ligand binding to E- and P-selectin analyzed using selectin/mannose-binding protein chimeras". *J. Biol. Chem.* 273: 6254-6261

- Tormo, J., Natarajan, K., Margulies, D.H., and Mariuzza, R.A. (1999). "Crystal structure of a lectin-like natural killer cell receptor bound to its MHC class I ligand". *Nature* 402: 623-631
- 5 Tsunetzawa, W., Sano, H., Sohma, H., McCormack, F.X., Voelker, D.R., and Kuroki, Y. (1998). "Site-directed mutagenesis of surfactant protein A reveals dissociation of lipid aggregation and lipid uptake by alveolar type II cells". *Biochim. Biophys. Acta* 1387: 433-446
- 10 Weis, W.I., Kahn, R., Fourme, R., Drickamer, K., and Hendrickson, W.A. (1991). "Structure of the calcium-dependent lectin domain from a rat mannose-binding protein determined by MAD phasing". *Science* 254: 1608-1615
- Weis, W.I., and Drickamer, K. (1996). "Structural basis of
15 lectin-carbohydrate recognition". *Annu. Rev. Biochem.* 65: 441-473
- Whitehorn, E.A., Tate, E., Yanofsky, S.D., Kochersperger, L., Davis A., Mortensen, R.B., Yonkovic, S., Bell, K., Dower, W.J., and Barrett, R.W. (1995). "A generic method
20 for expression and use of "tagged" soluble versions of cell surface receptors". *Bio/Technology* 13: 1215-1219
- Wragg, S. and Drickamer, K. (1999). "Identification of amino acid residues that determine pH dependence of ligand binding to the asialoglycoprotein receptor during endocytosis". *J. Biol. Chem.* 274: 35400-35406
25
- Zhang, H., Robison, B., Thorgaard, G.H., and Ristow, S.S. (2000). "Cloning, mapping and genomic organization of a fish C-type lectin gene from homozygous clones of rainbow trout (*Oncorhynchus Mykiss*)". *Biochim. et Biophys. Acta*
30 1494: 14-22

CLAIMS

1. A protein having the scaffold structure of C-type lectin-like domains (CTLD), said protein comprising a variant of a model CTLD wherein the α -helices and β -strands and connecting segments are conserved to such a degree that the scaffold structure of the CTLD is substantially maintained, while the loop region is altered by amino acid substitution, deletion, insertion or any combination thereof, with the proviso that said protein is not any of the known CTLD loop derivatives of C-type lectin-like proteins or C-type lectins listed in Table 2 in the description.
2. A protein according to claim 1 wherein the model CTLD is defined by having a 3D structure that conforms to the secondary-structure arrangement illustrated in Fig 1 characterized by the following main secondary structure elements:
- five β -strands and two α -helices sequentially appearing in the order β 1, α 1, α 2, β 2, β 3, β 4, and β 5, the β -strands being arranged in two anti-parallel β -sheets, one composed of β 1 and β 5, the other composed of β 2, β 3 and β 4,
- at least two disulfide bridges, one connecting α 1 and β 5 and one connecting β 3 and the polypeptide segment connecting β 4 and β 5, and
- a loop region consisting of two polypeptide segments, loop segment A (LSA) connecting β 2 and β 3 and comprising typically 15-70 or, less typically, 5-14 amino acid residues, and loop segment B (LSB) connecting β 3 and β 4 and comprising typically 5-12 or less typically, 2-4 amino acid residues.
3. A protein according to claim 1 wherein the model CTLD is defined by showing sequence similarity to a previously rec-

ognised member of the CTLD family as expressed by an amino acid sequence identity of at least 22 %, preferably at least 25 % and more preferably at least 30 %, and by containing the cysteine residues necessary for establishing the conserved two-disulfide bridge topology (i.e. Cys_I, Cys_{II}, Cys_{III} and Cys_{IV}), whereas the loop region and its flanking β -strand structural elements are identified by inspection of the sequence alignment with the collection of CTLDs shown in Fig. 1 providing identification of the sequence locations of the β 2- and β 3-strands with the further corroboration provided by comparison of these sequences with the four-residue consensus sequences, β 2cseq and β 3cseq, the β 4 strand segment being located typically at positions -6 to -2 and less typically at positions -5 to -2 relative to the conserved Cys_{III} residue and with the characteristic residues at positions -5 and -3 as elucidated from Table 1 and deducted in the description.

4. A protein according to any one of the preceding claims wherein up to 10, preferably up to 4, and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and/or β -strands and/or connecting segments of the model CTLD.

5. A protein according to any one of the preceding claims wherein changes of up to 4 residues are made in the β -strands of the model CTLD as a consequence of the introduction of recognition sites for one or more restriction endonucleases in the nucleotide sequence encoding the CTLD to facilitate the excision of part or all of the loop region and the insertion of an altered amino acid sequence instead while the scaffold structure of the CTLD is substantially maintained.

6. A protein according to any one of the preceding claims wherein the model CTLD is that of a tetranectin.

7. A protein according to claim 6 wherein the model CTLD is that of human tetranectin comprising amino acid residues 59 to 180 in the monomer native peptide sequence thereof.
8. A protein according to claim 6 wherein the model CTLD is that of murine tetranectin comprising amino acid residues 59 to 180 in the monomer native peptide sequence thereof.
9. A protein according to any one of the preceding claims which further comprises N-terminal and/or C-terminal extensions of the CTLD variant.
10. A protein according to claim 9 wherein said N-terminal and/or C-terminal extensions contain effector, enzyme, further binding and/or multimerising functions.
11. A protein according to claim 9 or 10 wherein said N-terminal and/or C-terminal extensions are the non-CTLD-portions of a native C-type lectin-like protein or C-type lectin or a "soluble" variant thereof lacking a functional transmembrane domain.
12. A protein according to any one of the preceding claims which is a multimer of a moiety comprising the CTLD variant.
13. A protein according to claim 12 which is derived from the native tetranectin trimer.
14. A protein according to claim 7 which is derived from the polypeptide htlec having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:13 by altering the amino acid sequence of the loop region.
15. A protein according to claim 7 which is derived from the polypeptide htCTLD having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:15 by altering the amino acid sequence of the loop region.

16. A protein according to claim 7 which is derived from the polypeptide hTN having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:09 by altering the amino acid sequence of the loop region.
- 5 17. A protein according to claim 7 which is derived from the polypeptide hTN3 having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:11 by altering the amino acid sequence of the loop region.
- 10 18. A protein according to claim 8 which is derived from the polypeptide mtlec having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:36 by altering the amino acid sequence of the loop region.
- 15 19. A protein according to claim 8 which is derived from the polypeptide mtCTLD having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:38 by altering the amino acid sequence of the loop region.
- 20 20. A combinatorial library of proteins having the scaffold structure of C-type lectin-like domains (CTLD), said proteins comprising variants of a model CTLD wherein the α -helices and β -strands are conserved to such a degree that the scaffold structure of the CTLD is substantially maintained, while the loop region or parts of the loop region of the CTLD is randomised with respect to amino acid sequence and/or number of amino acid residues.
- 25 21. A combinatorial library according to claim 20 wherein the model CTLD is defined by having a 3D structure that conforms to the secondary-structure arrangement illustrated in Fig. 1 characterised by the following main secondary structure elements:
- 30 five β -strands and two α -helices sequentially appearing in the order β 1, α 1, α 2, β 2, β 3, β 4, and β 5, the β -

strands being arranged in two anti-parallel β -sheets, one composed of $\beta 1$ and $\beta 5$, the other composed of $\beta 2$, $\beta 3$ and $\beta 4$,

5 at least two disulfide bridges, one connecting $\alpha 1$ and $\beta 5$ and one connecting $\beta 3$ and the polypeptide segment connecting $\beta 4$ and $\beta 5$, and

10 a loop region consisting of two polypeptide segments, loop segment A (LSA) connecting $\beta 2$ and $\beta 3$ and comprising typically 15-70 or, less typically, 5-14 amino acid residues, and loop segment B (LSB) connecting $\beta 3$ and $\beta 4$ and comprising typically 5-12 or less typically, 2-4 amino acid residues.

22. A combinatorial library according to claim 20 wherein the model CTLD is defined by showing sequence similarity to
15 a previously recognised member of the CTLD family as expressed by an amino acid sequence identity of at least 22 %, preferably at least 25 % and more preferably at least 30 %, and by containing the cysteine residues necessary for establishing the conserved two-disulfide bridge topology
20 (i.e. Cys_I, Cys_{II}, Cys_{III} and Cys_{IV}), whereas the loop region and its flanking β -strand structural elements are identified by inspection of the sequence alignment with the collection of CTLDs shown in Fig. 1 providing identification of the sequence locations of the $\beta 2$ - and $\beta 3$ -strands with
25 the further corroboration provided by comparison of these sequences with the four-residue consensus sequences, $\beta 2cseq$ and $\beta 3cseq$, the $\beta 4$ strand segment being located typically at positions -6 to -2 and less typically at positions -5 to -2 relative to the conserved Cys_{III} residue and with the
30 characteristic residues at positions -5 and -3 as elucidated from Table 1 and deducted in the description.

23. A combinatorial library according to any one of claims 20-22 of proteins comprising CTLD variants wherein up to

10, preferably up to 4, and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and β -strands and connecting segments of the model CTLD.

5 24. A combinatorial library according to any one of claims 20-23 of proteins comprising CTLD variants wherein changes of up to 4 residues are made in the model CTLD as a consequence of the introduction of recognition sites for one or more restriction endonucleases in the nucleotide sequence
10 encoding the CTLD to facilitate the excision of part or all of a DNA segment encoding the loop region and the insertion of members of an ensemble of DNA fragments that collectively encode a randomised amino acid sequence instead while the scaffold structure of the CTLD is substantially
15 maintained.

25. A combinatorial library according to any one of claims 20-24 of proteins wherein the model CTLD is that of a tetranectin.

20 26. A combinatorial library according to claim 25 of proteins wherein the model CTLD is that of human tetranectin comprising amino acid residues 59 to 180 in the monomer native peptide sequence thereof.

25 27. A combinatorial library according to claim 25 of proteins wherein the model CTLD is that of murine tetranectin comprising amino acid residues 59 to 180 in the monomer native peptide sequence thereof.

28. A combinatorial library according to any one of claims 20-27 of proteins which further comprise N-terminal and/or C-terminal extensions of the CTLD-variant.

30 29. A combinatorial library according to claim 28 of proteins wherein said N-terminal and/or C-terminal extensions

contain effector, enzyme, further binding and/or multimerising functions.

30. A combinatorial library according to claim 28 or 29 of proteins wherein said N-terminal and/or C-terminal extensions are the non-CTLD-portions of a native C-type lectin-like protein or C-type lectin or a "soluble" variant thereof lacking a functional transmembrane domain.
31. A combinatorial library according to any one of claims 20-30 of proteins which are multimers of a moiety comprising the CTLD variant.
32. A combinatorial library according to claim 31 of proteins which are derived from the native tetranectin trimer.
33. A combinatorial library according to claim 27 of proteins which are derived from the peptide htlec having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:13 by randomising the loop region with respect to amino acid sequence and/or number of amino acid residues.
34. A combinatorial library according to claim 27 of proteins which are derived from the peptide htCTLD having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:15 by randomising the loop region with respect to amino acid sequence and/or number of amino acid residues.
35. A combinatorial library according to claim 27 of proteins which are derived from the peptide hTN having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:09 by randomising the loop region with respect to amino acid sequence and/or number of amino acid residues.

36. A combinatorial library according to claim 27 of proteins which are derived from the peptide hTN3 having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:11 by randomising the loop region with respect
5 to amino acid sequence and/or number of amino acid residues.
37. A combinatorial library according to claim 28 of proteins which are derived from the peptide mtlec having the amino acid sequence from position 5 Glu to position 185 Val
10 in SEQ IN NO:36 by randomising the loop region with respect to amino acid sequence and/or number of amino acid residues.
38. A combinatorial library according to claim 28 of proteins which are derived from the peptide mtCTLD having the amino acid sequence from position 5 Ala to position 141 Val
15 in SEQ IN NO:38 by randomising the loop region with respect to amino acid sequence and/or number of amino acid residues.
39. A derivative of a native tetranectin wherein up to 10, preferably up to 4, and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and/or β -strands and/or connecting segments of its CTLD with the proviso that said derivative is not any of the known CTLD derivatives of human tetranectin (hTN)
20 listed in Table 2 in the description.
40. A derivative of human tetranectin, termed htlec, having the amino acid sequence from position 5 Glu to position 185 Val in SEQ IN NO:13.
41. A derivative of human tetranectin, termed htCTLD, having the amino acid sequence from position 5 Ala to position
30 141 Val in SEQ IN NO:15.

42. A derivative of human tetranectin, termed PhTN, having the amino acid sequence given in SEQ IN NO:09.
43. A derivative of human tetranectin, termed PhTN3, having the amino acid sequence given in SEQ IN NO:11.
- 5 44. A derivative of human tetranectin, termed Phtlec, having the amino acid sequence given in SEQ IN NO:13.
45. A derivative of human tetranectin, termed PhtCTLD, having the amino acid sequence given in SEQ IN NO:15.
46. A derivative of human tetranectin, termed FX-htlec,
10 having the amino acid sequence given in SEQ IN NO:02.
47. A derivative of human tetranectin, termed FX-htCTLD, having the amino acid sequence given in SEQ IN NO:04.
48. A derivative of murine tetranectin, termed mtlec, having the amino acid sequence from position 5 Glu to position
15 185 Val in SEQ IN NO:36.
49. A derivative of murine tetranectin, termed mtCTLD, having the amino acid sequence from position 5 Ala to position 141 Val in SEQ IN NO:38.
50. A derivative of murine tetranectin, termed Pmtlec, having
20 the amino acid sequence given in SEQ IN NO:36.
51. A derivative of murine tetranectin, termed PmtCTLD, having the amino acid sequence given in SEQ IN NO:38.
52. A derivative of murine tetranectin, termed FX-mtlec, having the amino acid sequence given in SEQ IN NO:29.
- 25 53. A derivative of murine tetranectin, termed FX-mtCTLD, having the amino acid sequence given in SEQ IN NO:31.

54. Nucleic acid comprising a nucleotide sequence encoding a htlec insert as shown from nucleotide 20 to nucleotide 562 in SEQ ID NO:12.
55. Nucleic acid comprising a nucleotide sequence encoding
5 a htCTLD insert as shown from nucleotide 20 to nucleotide 430 in SEQ ID NO:14.
56. Nucleic acid comprising a nucleotide sequence encoding a mtlec insert as shown from nucleotide 20 to nucleotide 562 in SEQ ID NO:35.
- 10 57. Nucleic acid comprising a nucleotide sequence encoding a mtCTLD insert as shown from nucleotide 20 to nucleotide 430 in SEQ ID NO:37.
58. Nucleic acid comprising any nucleotide sequence encoding a protein according to any one of claims 1-19.
- 15 59. A library of nucleic acids encoding proteins of a combinatorial library according to any one of claims 20-38, in which the members of the ensemble of nucleic acids, that collectively constitute said library of nucleic acids, are able to be expressed in a display system, which provides
20 for a logical, physical or chemical link between entities displaying phenotypes representing properties of the displayed expression products and their corresponding genotypes.
60. A library of nucleic acids according to claim 49,
25 wherein the display system is selected from
- (I) a phage display system such as
 - (1) a filamentous phage fd in which the library of nucleic acids is inserted into
 - (a) a phagemid vector,
 - (b) the viral genome of a phage
- 30

- (c) purified viral nucleic acid in purified single- or double-stranded form, or
- (2) a phage lambda in which the library is inserted into
- (a) purified phage lambda DNA, or
- (b) the nucleic acid in lambda phage particles; or
- (II) a viral display system in which the library of nucleic acids is inserted into the viral nucleic acid of a eukaryotic virus such as baculovirus; or
- (III) a cell-based display system in which the library of nucleic acids is inserted into, or adjoined to, a nucleic acid carrier able to integrate either into the host genome or into an extrachromosomal element able to maintain and express itself within the cell and suitable for cell-surface display on the surface of
- (a) bacterial cells,
- (b) yeast cells, or
- (c) mammalian cells; or
- (IV) a nucleic acid entity suitable for ribosome linked display into which the library of nucleic acid is inserted; or
- (V) a plasmid suitable for plasmid linked display into which the library of nucleic acid is inserted.

61. A library of nucleic acids according to claim 60 wherein said phagemid vector is the vector "pCANTAB 5 E" supplied by Amersham Pharmacia Biotech (code no. 27-9401-01) for use with their "Recombinant Phage Antibody System".

62. A method of preparing a protein according to any one of claims 1-19, wherein the protein comprises at least one or

more, identical or not identical, CTLD domains with novel loop-region sequences which has (have) been isolated from one or more CTLD libraries by screening or selection.

63. A method of preparing a protein according to claim 62,
5 wherein at least one CTLD domain has been further modified by mutagenesis.

64. A method of preparing a protein according to claim 62 or 63, wherein the protein containing at least one CTLD domain is assembled from two or more components by chemical
10 or enzymatic coupling or crosslinking.

65. A method of preparing a combinatorial library according to any one of claims 20-38 comprising the following steps:

- 1) inserting nucleic acid encoding a protein comprising a model CTLD into a suitable vector,
- 15 2) if necessary, introducing restriction endonuclease recognition sites by site directed mutagenesis, said recognition sites being properly located in the sequence at or close to the ends of the sequence encoding the loop region of the CTLD or part thereof,
- 20 3) excising the DNA fragment encoding the loop region or part thereof by use of the proper restriction endonucleases,
- 4) ligating mixtures of DNA fragments into the restricted vector, and
- 25 5) inducing the vector to express randomised proteins having the scaffold structure of CTLDs in a suitable medium.

66. A method of constructing a tetranectin derivative adapted for the preparation of a combinatorial library according to any one of claims 20-38, wherein the nucleic
30 acid encoding the tetranectin derivative has been modified to generate endonuclease restriction sites within nucleic acid segments encoding $\beta 2$, $\beta 3$ or $\beta 4$, or up to 30 nucleo-

tides upstream or downstream in the sequence from any nucleotide which belongs to a nucleic acid segment encoding $\beta 2$, $\beta 3$ or $\beta 4$.

- 5 67. The use of a nucleotide sequence encoding a tetranectin, or a derivative thereof wherein the scaffold structure of its CTLD is substantially maintained, for preparing a library of nucleotide sequences encoding related proteins by randomising part or all of the nucleic acid sequence encoding the loop region of its CTLD.
- 10 68. The use according to claim 67 wherein the nucleotide sequence encodes a mammalian tetranectin,
69. The use according to claim 67 wherein the nucleotide sequence encodes human or murine tetranectin.
- 15 70. The use according to claim 67 wherein the nucleotide sequence encodes a derivative of a native tetranectin wherein up to 10, preferably up to 4, and more preferably 1 or 2, amino acid residues are substituted, deleted or inserted in the α -helices and β -strands and connecting segments of its CTLD.
- 20 71. The use according to claim 67 wherein the nucleotide sequence encodes a derivative of human tetranectin, termed htlec, as shown from nucleotide 20 to nucleotide 562 in SEQ ID NO:12.
- 25 72. The use according to claim 67 wherein the nucleotide sequence encodes a derivative of human tetranectin, termed htCTLD, as shown from nucleotide 20 to nucleotide 430 in SEQ ID NO:14.
73. The use according to claim 67 wherein the nucleotide sequence encodes a derivative of murine tetranectin, termed

mtlec, as shown from nucleotide 20 to nucleotide 562 in SEQ ID NO:35.

74. The use according to claim 67 wherein the nucleotide sequence encodes a derivative of murine tetranectin, termed
5 mtCTLD, as shown from nucleotide 20 to nucleotide 430 in SEQ ID NO:37.

75. A method of screening a combinatorial library according to any one of claims 20-38 for binding to a specific target which comprises the following steps:

- 10 1) expressing a nucleic acids library according to any one of claims 59-61 to display the library of proteins in the display system;
- 2) contacting the collection of entities displayed with a suitably tagged target substance for which isolation
15 of a CTLD-derived exhibiting affinity for said target substance is desired;
- 3) harvesting subpopulations of the entities displayed that exhibit affinity for said target substance by means of affinity-based selective extractions, utiliz-
20 ing the tag to which said target substance is conjugated or physically attached or adhering to as a vehicle or means of affinity purification, a procedure commonly referred to in the field as "affinity panning", followed by re-amplification of the sub-
25 library;
- 4) isolating progressively better binders by repeated rounds of panning and re-amplification until a suitably small number of good candidate binders is obtained; and,
- 30 5) if desired, isolating each of the good candidates as an individual clone and subjecting it to ordinary functional and structural characterisation in preparation for final selection of one or more preferred product clones.

76. A method of reformatting a protein according to any one of claims 1-19 or selected from a combinatorial library according to any one of claims 20-38 and containing a CTLD variant exhibiting desired binding properties, in a desired
5 alternative species-compatible framework by excising the nucleic acid fragment encoding the loop region-substituting polypeptide and any required single framework mutations from the nucleic acid encoding said protein using PCR technology, site directed mutagenesis or restriction enzyme di-
10 gestion and inserting said nucleic acid fragment into the appropriate location(s) in a display- or protein expression vector that harbours a nucleic acid sequence encoding the desired alternative CTLD framework.

1/34

[illegible]

Fig. 1




Fig. 2

htlec
mtlec

E P P T Q K P K K I V N A K K D V V N T K M F E L K S R L 30
GAGCCACCAACCCAGAACGATGTAAATGCAAGAAGATGTGTGAACACAAAGATGTTGAGGAGCTCAAGAGCCGCTCG 90
GAGTACCCCACTCCCAAGGCCAAGAGGCTGCAATGCCAAGAAAGATTGGTGAGCTCAAAGATGTTGAGGAGCTCAAGAACAGGATG
E S P T P K A K K A A N A K K D L V S S K M F E E L K N R M

91
D T L A Q E V A L L K E Q Q A L Q T V V L K G T K V H M K V 60
GACACCTGGCCAGGAGGTGGCCCTGCTGAAGAGCAGAGCCCTGCAGACGGTCTGCTGAAGGGGACCAAGGTGCACATCAAAAGTC 180
GATGTCTGGCCCAAGGAGGTGGCCCTGCTGAAGAGAGCAGGCCCTTACAGACTGTGTCTCTGAAGGGCACCAAGGTGAACCTTGAAGGTC
D V L A Q E V A L L K E K Q A L Q T V V L K G T K V N L K V

181
F L A F T Q T K' T F H E A S E D C I S R G G T L S T P Q T G 90
TTTCTGGCCTTACCCAGACGAAGACCTTCCACGAGGCCAGCAGGACTGCATCTCGCGGGGGCACCTTGAGCACCCCTCAGACTGGC 270
CTCCTGGCCTTACCCAAACCGAAGACCTTCCATGAGCGAGCAGGACTGCATCTCGAAGGGGCACGCTGGGCACCCCGCAGTCAGAG
L L A F T Q P K T F H E A S E D C I S Q G G T L G T P P Q S E

271
S E N D A L Y E Y L R Q S V G N E A E I W L G L N D M A A E 120
TCGGAGAACGACGCCCTGTATGAGTACCTGCGCCAGAGCCTGGGCAACGAGGCCGAGATCTGGCTGGGCCCTCAACGACATGGCGCCGAG 360
CTAGAGAACGAGGCGCTGTTGAGTACGCGCCACAGCCTGGGCAACGATGCGGAGATCTGGCTGGGCCCTCAACGACATGGCCCGCGAA
L E N E A L F E Y A R H S V G N D A E I W L G L N D M A A E

361
G T W V D M T G T R I A Y K N W E T E I T A Q P D G G K T E 150
GGCAGCTGGGTGGACATGACCGGTACCCGTCATCGCTTACAAAGAACTGGGAGACTGAGATCACCGCGCAACCCGATGGCGGCAAGACCGAG 450
GGCGCTGGGTGGACATGACCGGTACCCCTGCGCTTACAAAGAACTGGGAGACGGAGATCACGACGCAACCCGACGGCGGCAAGCCGAG
G A W V D M T G T L L A Y K N W E T E I T T Q P D G G K A E

451
N C A V L S G A A N G K W F D K R C R D Q L P Y I C Q F G I V *
AACTGCGCGGCTGTGTCAGGCGCGCCAAACGGCAAGTGGTTCGACAAAGCGTGGCCGATCAATTGCCCTACATCTGCCAGTTTCGGGATCGTGTA
AACTGCGCGGCTGTGTCAGGCGAGCCAAACGGCAAGTGGTTCGACAAAGCGATGCCCGGATCAATTGCCCTACATCTGCCAGTTTGCCCATGTGTGA
N C A A L S G A A N G K W F D K R C R D Q L P Y I C Q F A I V *
Mun I

3/34

Fig. 3

4/34

h1CTLD 30
 A L Q T V V L K G T K V H M K V F L A F T Q T K T F H E A S
 G C C T G C A G A C G G T C G T C C T G A A G G G A C C A A G G T G C A C A T G A A A G T C T T T C T G G C C T T C A C C C A G A C G A A G A C C T T C C A G A G G C C A G C
 m1CTLD 90
 G C C T T A C A G A C T G T G G T C C T G A A G G G C A C C A A G G T G A A G G T C C T C C T G G C C T T C A C C C A A C C G A A G A C C T T C C A T G A G G C G A G C
 A L Q T V V L K G T K V N L K V L L A F T Q P K T F H E A S

 91 60
 E D C I S R G G T L S T P Q T G S E N D A L Y E Y L R Q S V
 G A G G A C T G C A T C T C G C G G G G C A C C C T G A G A C A C C C T C A G A C T G G C T G G A G A A C G A C G C C C T G T A T G A T A C C T G C G C C A G A G C G T G
 G A G G A C T G C A T C T C G C A A G G G G C A C G T G G G C A C C C C G C A G T C A G A G C T A G A G A A C G A G G C G C T G T T C G A G T A C G C G C C C A C A G C G T G
 E D C I S Q G G T L G T P Q S E L E N E A L F E Y A R H S V

 181 90
 G N E A E I W L G L N D M A A E G T W V D M T G T R I A Y K
 G G C A A C G A G C C G A G A T C T G G C T G G G C C T C A A C G A C A T G G C G C G A G G C A C C T G G G T G G A C A T G A C C G G T A C C C G C A T C G C C T A C A A G
 G G C A A C G A T G C G G A G A T C T G G C T G G G C C T C A A C G A C A T G G C C G C G A A G G C G C C T G G G T G G A C A T G A C C G G T A C C C T C C T G G C C T A C A A G
 G N D A E I W L G L N D M A A E G A W V D M T G T L L A Y K
 Bgl II

 271 120
 N W E T E I T A Q P D G G K T E N C A V L S G A A N G K W F
 A A C T G G G A G A C T G A G A T C A C C G C G A A C C C G A T G G C G G C A A G A C C G A G A A C T G C G C G G T C C T G T C A G G C G C G G C C A A C G G C A A G T G G T T C
 A A C T G G G A G A C G G A G A T C A C G A C G A A C C C G A C G G C G G C A A G C C G A G A A C T G C G C G C C C T G T C T G G C G C A G C C A A C G G C A A G T G G T T C
 N W E T E I T T Q P D G G K A E N C A A L S G A A N G K W F
 Kpn I

 361
 D K R C R D Q L P Y I C Q F G I V *
 G A C A A G C G T G C C G G A T C A A T I G C C C T A C A T C T G C C A G T T C G G G A T C G T G T A G
 G A C A A G C G A T G C C G G A T C A A T T G C C C T A C A T C T G C C A G T T G C C A T T G T G T A G
 D K R C R D Q L P Y I C Q F A I V *
 Mun I

Fig. 4

pT7H6FX-htlec

G S I E G R G E P P T -- V V -- K V E A E I W -- G T R -- Q L P Y I C Q F G I V *

G S I E G R G E P T -- V V -- K V E A E I W -- G T R -- Q L P Y I C Q F G I V *
GATCCATCGAGGTAGGGCGAGCCACCAACC---GTCGTC---AAAGTC---GAGGCCGAGATCTGG---GGTACCCGC---CAATTGCCCTCATCTGCCAGTTCCGGGATCGGTGA

Bgl II	Kpn I	Mun I
--------	-------	-------


M G S H H H H H
 CATATGGGATCGCATCACCATCACCATCAGG --- AGCTTGAATTC
 Bam HI Hind III

M G S H H H H H
CATATGGGATCGCATCACCATCACCATCAG

Bam Hi

- - AGCTTGAATTC

III. PUIH



Ori

pT7H6FX-htlec

Amp^r

Fig. 5

6/34

FX-htlec

```

1  G S I E G R G E P P P T Q K P K K I V N A K K D V V N T K M F
31 E E L K S R R L D T L A Q E V A L L K H E Q Q A A L D Q T V V R L K G
61 T K V L S T P H M Q K T G L V F L F H E Q A S S V G N D Q C I S R G T
91 L S T P M Q A T G E N C A W A N G K Y W F D E E I I W L A L
121 L N D G G K F Q T E N V A V L S G A A N G K F E E T C R E I I T Q
151 P D G G K F Q T E N V A V L S G A A N G K F E E T C R E I I T Q
181 Y I C Q F G I V A V L S G A A N G K F E E T C R E I I T Q

```

Fig. 6

7/34

pT7H6FX-htCTL D

G S I E G R A L Q T V V -- K V -- E A E I W -- G T R -- Q L P Y I C Q F G I V *
 GATCCATCGAGGGTAGGGCCCTGCAGACGGTCGTC--AAAGTC--GAGGCCGAGATCTGG--GGTACCCGC--CAATTGCCCTACATCTGCCAGTTCGGGATCGTGTA

Mun I

Kpn I

Bgl II

M G S H H H H H H

CATATGGGATCGCATCACCATCACCACG

Bam HI

AGCTTGAATTC

Hind III

T7-phage
promoter

Óri

pT7H6FX-htCTL D

Amp^r

Fig.7

FX-htCTL D

1 G S I E G R A L Q T V V L K G T L S K V H M K V F L A F T Q T K
31 T F L R Q S E A C I S R G G T L S P Q T G N D A D Q Y E
61 Y L R Q S E A C I S R G G T L S P Q T G N D A D Q Y E
91 T R I A Y K N W E E T C R D Q L P Y I C G Q F G I
121 A N G K W F D K R C R D Q L P Y I C G Q F G I

Fig.8

9/34

pPhTN

P A M A E P P T -- V C -- K C -- E A E I W -- G A R -- Q L P Y I C Q F G I V A
CGGCCATGGCCGAGCCACCAACC--GTCTGC--AAATGC--GAGGCCGAGATCTGG--GGCGCCCGC--CAGCTGCCCTACATCTGCCAGTTCGGGATCGTGGC

Y A A Q
TATGCGGCCCAGC Sfi I
A A G A
GGCCGCAGGTGCG Not I




Fig.9

10/34

PhTN

```

1 P A M A E P T Q K P K K I V N A K K D V V T N T K M F E E L
31 K S R L D T A Q E V A L L K E Q Q A L C R L G L T K S V
61 H M K C F S E T L A F T Q T Y E K S E D C I S V R G T L S T
91 P Q T G S E N W C A V D A L M S G A N R I Q A S V K N D E E
121 M A A E N C A V D A L M S G A N R I Q A S V K N D E E
151 G K T E N C A V D A L M S G A N R I Q A S V K N D E E
181 Q F G I V A A A A A A A A A A A A A A A A A A A A A A

```

Fig.10

11/34

pPhTN3

P A M A A L Q T V C -- K C -- D A E I W -- G A R -- Q L P Y I C Q F G I V A
CGGCCATGGCCGCCCTGCAGACGGTCTGC--AAATGC--GAGGCCGAGATCTGG--GGCGCCCGC--CAGCTGCCCTACATCTGCCAGTTCCGGGATTGTGGC

Y A A Q
TATGGGCCCCAGC Sfi I
A A G A
GGCCGCAGGTGCG Not I




Fig.11

12/34

PhTN3

1 P A M A A L D C I S R I I T A Q L P Y I C Q F G I V A A
31 H E A S V G N W K F D K R C R D Q L P Y I C Q F G I V A A
61 R Q S Y K N W K F D K R C R D Q L P Y I C Q F G I V A A
91 I A Y K N W K F D K R C R D Q L P Y I C Q F G I V A A
121 G K W F D K R C R D Q L P Y I C Q F G I V A A

Fig.12

pPhtlec

P A M A E P P T -- V -- V -- K V -- E A E I W -- G T R -- Q L P Y I C Q F G I V A
 CGGCCATGGCCGAGCCACCAACC--GTCGTC--AAAGTC--GAGGCCGAGATCTGG--GGTACCCCGC--CAATTGCCCTACATCTGCCAGTTTCGGGATCGTGGC

Bgl II Kpn I Mun I

Y A A Q
 TATGGGGCCCAGC Sfi I
 A A G A
 GGCCGCAGGTGCG Not I




Fig.13

14/34

Phtlec

```

1 P A M A E D F F S S E G T C A A
31 K S R K V G S E E N V A A
61 H M K T G A E E N V A A
91 P Q A A T E E N V A A
121 M A A E E N V A A
151 G K T E E N V A A
181 Q F G I V A A

T Q Q T A F D V V A
K K E Q T L M S
P V T Y T G A
K A K E G A
K L T Y T A N
I L F L R I G
V K H E Q A K
N E E A S Y W
A Q Q A S V K F
K K A A D N W K
V D L Q C E E R
V T Q I A E R
N V S E E R
T V R I I D
K L G W T Q
M K G L A L
F G T G Q P
E T L L P Y
E K S N D I
L V T D G C

```

Fig.14

pPhitCTLD

P A M A A L Q T V V -- K V -- E A E I W -- G T R -- Q L P Y I C Q F G I V A
CGCCCATGGCCCGCCCTGCAGACGGTCGTC--AAAGTC--GAGCCGAGATCTGG--GGTACCCGC--CAATTGCCCTACATCTGCCAGTTCGGGATGTGGC

Bgl II

Kpn I

Mun I

Y A A Q

TATGCGCCCGC

Sfi I

A A G A

GGCCGCAGGTGCG

Not I




Fig.15

16/34

Ph_tCTL_D

1 P A M A A L Q T I S S R G I I T A Q P Y I C Q F G I V A A
31 H E A S E D C E A E I I T A Q P Y I C Q F G I V A A
61 R Q S V G N E E T E R C R D Q L L P Y I C Q F G I V A A
91 I A Y K N W E E T E R C R D Q L L P Y I C Q F G I V A A
121 G K W F D K R C R D Q L L P Y I C Q F G I V A A

Fig.16

pUC-mtlec




Fig.17

pT7H6FX-mtlec

S I Q G R G E S P T -- V -- K V -- D A E I W -- G T L -- Q L P Y I C Q F A I V *
 TCCATCCAGGGTCGGCGGAGTCACCCACT--GTGGTC--AAGGTC--GATGCGGAGATCTGG--GGTACCCCTC--CAATTGCCCTACATCTGCCAGTTTGCCCATTTGTGTA

Bgl II Kpn I Mun I

M G S H H H H H H

CATATGGGATCGCATCACCATCACCATCAGC

Bam HI

AGCTTGAATTC

Hind III

T7-phage promoter

Ori

pT7H6FX-mtlec

Amp^r

Fig.18

19/34

FX-mtlec

```

1  G S I Q G R G E S P T P K A K A A N A K K D L V S S K M F
31 E E L K N L Q S E R M D V L A F A Q E V A L K E K Q A L Q T V V L K G
61 T K V N L Q S E R M D V L A F A Q E V A L K E K Q A L Q T V V L K G
91 L G T P M A A E N C A W A L M T G T L A Y S V K N D K R I I T W L G
121 L N D M A A E N C A W A L M T G T L A Y S V K N D K R I I T W L G
151 P D G G K A E N C A W A L M T G T L A Y S V K N D K R I I T W L G
181 Y I C Q F A I V

```

Fig.19




Fig.20

EX-mtCTLD

1 G S I Q G R A L Q T V V L K G T K V N L K V L L A F T Q P K
31 T F H E A S E D C I S Q G T L G T L G T P Q S E L E N E A L F F E
61 Y A R H S V G N D A E I W L T T Q P D G G K A E N C A A L S G A
91 T L A Y K N W E R C R D Q L P Y I C Q F A I V
121 A N G K W F D K R C R D Q L P Y I C Q F A I V

Fig. 21

pPmtlec

P A M A E S P T -- V -- V -- K V -- D A E I W -- G T L -- Q L P Y I C Q F A I V A
CGGCCATGGCCGAGTCACCAACC--GTGGTC--AAGGTC--GATGGGAGATCTGG--GGTACCCCTC--CAATTGCCCTACATCTGCCAGTTTGCCCATTTGTGGC

Bgl II Kpn I Mun I

Y A A Q
TATGCGGCCCAGC Sfi I
A A G A
GGCCGCAGGTGCG Not I




Fig.22

23/34

Pmtlec

1 P A M A E S P T P K A K K A A N A K K D L V S S V S K M F E E L
31 K N R M D V L A F T Q E V A L L K H E K Q A L Q T V V Q L K G T K V
61 N L K V L L E N E A L F E Y A R H E A S E D C I S Q G T L G T K V
91 P Q S E L E G A W V D M T G T L A Y K N D E R A E I W T T L P Y I
121 M A A E E N C A A L S G A N G K R C D I D Q L L P Y I C
151 G K A E N C A A L S G A N G K R C D I D Q L L P Y I C
181 Q F A I V A A A

Fig.23

pPmtCTLD

P A M A A L Q T V V -- K V -- D A E I W -- G T L -- Q L P Y I C Q F A I V A
 CGGCCATGGCCGCTTACAGACTGTGGTC--AAGGTC--GATGCGGAGATCTGG--GGTACCCCTC--CAATTGCCCTACATCTGCCAGTTTGCCCATTTGTGGC

Bgl II Kpn I Mun I

Y A A Q
 TATGCGGCCAGC Sfi I
 A A G A
 GGCCGCAGGTGCG Not I




Fig.24

25/34



PmtCTL

1 P A M A A L Q T V V L K G T K V N L K V L L A F T Q P K T F
31 H E A S E D C I S Q G G T L G T P Q S E L L E N E A L F E Y A
61 R H S V G N D A E I T T Q P D G G K A A E V D M S G T L
91 L A Y K N W E T E I T T Q P D G G K A A E V D M S G T L
121 G K W F D K R C R D Q L P Y I C Q F A I V A A A

Fig.25



26/34

Figure 26



27/34

Figure 27



28/34

Figure 28





Figure 29



29/34

PrMBP

1 P A M A N K L H A F S M G K K S G K K F F V T N H E R M P
31 F S K V K A L C S E L T D E V T A I G Q E
61 V A K T S A F L G I T H G S G E D C V
91 S N W K K D E P N D H C E F P A A
121 S C Q A S H T A V C E F P A A

Fig.30

pPrMBP

P A M A N K L H -----E F P A A
CGCCCATGGCCCAACAAGTTGCAT-----GAGTCCCGAGCCGC

Y A A Q
TATGCGGCCCAGC -----A A G A
Sfi I -----GGCCGCAGGTGCG
Not I




Fig.31

PhSP-D

1 P A M A K K V E L F P N G G Q S V G E K I F K T A G F V K P
31 F T E A Q L L C T Q A G G Q Q L A S P R S A A E N A A L Q Q
61 L V V A K N W A P G E L P V C E F A A
91 L V Y S N W A P G E L P V C E F A A
121 N D R A C G E K R L V C E F A A

Fig. 32

32/34

pPhSP-D

P A M A K K V E ---V C E F A
 C G G C C A T G G C C A A G A A G T T G A G ---G T C T G C G A G T T C G C

Y A A Q
 T A T G C G G C C C A G C
 Sfi I

A A G A
 G G C C G C A G G T G C G
 Not I






Fig.33

Figure 34



34/34

Figure 35



SEQUENCE LISTING

<110> Borean Pharma A/S

<120> Combinatorial libraries of proteins having the scaffold structure of C-type lectin-like domains

<130> P200001100 WO JNy

<140>

<141>

<150> DK PA 2000 01872

<151> 2000-12-13

<160> 91

<170> PatentIn Ver. 2.1

<210> 1

<211> 571

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(564)

<223> FX-htlec encoding insert

<400> 1

gga tcc atc gag ggt agg ggc gag cca cca acc cag aag ccc aag aag	48
Gly Ser Ile Glu Gly Arg Gly Glu Pro Pro Thr Gln Lys Pro Lys Lys	
1 5 10 15	

att gta aat gcc aag aaa gat gtt gtg aac aca aag atg ttt gag gag	96
Ile Val Asn Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu	
20 25 30	

ctc aag agc cgt ctg gac acc ctg gcc cag gag gtg gcc ctg ctg aag	144
Leu Lys Ser Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys	
35 40 45	

gag cag cag gcc ctg cag acg gtc gtc ctg aag ggg acc aag gtg cac	192
Glu Gln Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val His	
50 55 60	

atg aaa gtc ttt ctg gcc ttc acc cag acg aag acc ttc cac gag gcc	240
Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala	
65 70 75 80	

agc gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc cct cag act	288
Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr	
85 90 95	

ggc tcg gag aac gac gcc ctg tat gag tac ctg cgc cag agc gtg ggc	336
Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly	
100 105 110	

aac gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg gcc gag ggc	384
Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly	
115 120 125	

acc tgg gtg gac atg acc ggt acc cgc atc gcc tac aag aac tgg gag 432
 Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys Asn Trp Glu
 130 135 140

act gag atc acc gcg caa ccc gat ggc ggc aag acc gag aac tgc gcg 480
 Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala
 145 150 155 160

gtc ctg tca ggc gcg gcc aac ggc aag tgg ttc gac aag cgc tgc cgc 528
 Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg
 165 170 175

gat caa ttg ccc tac atc tgc cag ttc ggg atc gtg taagctt 571
 Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val
 180 185

<210> 2

<211> 188

<212> PRT

<213> Homo sapiens

<400> 2

Gly Ser Ile Glu Gly Arg Gly Glu Pro Pro Thr Gln Lys Pro Lys Lys
 1 5 10 15

Ile Val Asn Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu
 20 25 30

Leu Lys Ser Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys
 35 40 45

Glu Gln Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val His
 50 55 60

Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala
 65 70 75 80

Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr
 85 90 95

Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly
 100 105 110

Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly
 115 120 125

Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys Asn Trp Glu
 130 135 140

Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala
 145 150 155 160

Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg
 165 170 175

Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val
 180 185

<210> 3
 <211> 436
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(429)
 <223> FX-htCTLD encoding insert

<400> 3
 gga tcc atc gag ggt agg gcc ctg cag acg gtc gtc ctg aag ggg acc 48
 Gly Ser Ile Glu Gly Arg Ala Leu Gln Thr Val Val Leu Lys Gly Thr
 1 5 10 15
 aag gtg cac atg aaa gtc ttt ctg gcc ttc acc cag acg aag acc ttc 96
 Lys Val His Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe
 20 25 30
 cac gag gcc agc gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc 144
 His Glu Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr
 35 40 45
 cct cag act ggc tcg gag aac gac gcc ctg tat gag tac ctg cgc cag 192
 Pro Gln Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln
 50 55 60
 agc gtg ggc aac gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg 240
 Ser Val Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
 65 70 75 80
 gcc gag ggc acc tgg gtg gac atg acc ggt acc cgc atc gcc tac aag 288
 Ala Glu Gly Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys
 85 90 95
 aac tgg gag act gag atc acc gcg caa ccc gat ggc ggc aag acc gag 336
 Asn Trp Glu Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu
 100 105 110
 aac tgc gcg gtc ctg tca ggc gcg gcc aac ggc aag tgg ttc gac aag 384
 Asn Cys Ala Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
 115 120 125
 cgc tgc cgc gat caa ttg ccc tac atc tgc cag ttc ggg atc gtg 429
 Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val
 130 135 140
 taagctt 436

<210> 4
 <211> 143
 <212> PRT
 <213> Homo sapiens

<400> 4
 Gly Ser Ile Glu Gly Arg Ala Leu Gln Thr Val Val Leu Lys Gly Thr
 1 5 10 15
 Lys Val His Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe
 20 25 30

His Glu Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr
 35 40 45
 Pro Gln Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln
 50 55 60
 Ser Val Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
 65 70 75 80
 Ala Glu Gly Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys
 85 90 95
 Asn Trp Glu Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu
 100 105 110
 Asn Cys Ala Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
 115 120 125
 Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val
 130 135 140

<210> 5
 <211> 47
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 5
 cggctgagcg gccagccgg ccatggccga gccaccaacc cagaagc

47

<210> 6
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 6
 cctgcggccg ccacgatccc gaactgg

27

<210> 7
 <211> 43
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 7
 cggctgagcg gccagccgg ccatggccgc cctgcagacg.gtc

43

<210> 8
 <211> 570
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (8)..(565)
 <223> PhTN encoding insert

<400> 8

```

ggcccag ccg gcc atg gcc gag cca cca acc cag aag ccc aag aag att 49
      Pro Ala Met Ala Glu Pro Pro Thr Gln Lys Pro Lys Lys Ile
        1              5              10

gta aat gcc aag aaa gat gtt gtg aac aca aag atg ttt gag gag ctc 97
Val Asn Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu Leu
  15              20              25              30

aag agc cgt ctg gac acc ctg gcc cag gag gtg gcc ctg ctg aag gag 145
Lys Ser Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys Glu
              35              40              45

cag cag gcc ctg cag acg gtc tgc ctg aag ggg acc aag gtg cac atg 193
Gln Gln Ala Leu Gln Thr Val Cys Leu Lys Gly Thr Lys Val His Met
              50              55              60

aaa tgc ttt ctg gcc ttc acc cag acg aag acc ttc cac gag gcc agc 241
Lys Cys Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala Ser
        65              70              75

gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc cct cag act ggc 289
Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr Gly
      80              85              90

tcg gag aac gac gcc ctg tat gag tac ctg cgc cag agc gtg ggc aac 337
Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly Asn
      95              100              105              110

gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg gcc gag ggc acc 385
Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Thr
              115              120              125

tgg gtg gac atg acc ggc gcc cgc atc gcc tac aag aac tgg gag act 433
Trp Val Asp Met Thr Gly Ala Arg Ile Ala Tyr Lys Asn Trp Glu Thr
              130              135              140

gag atc acc gcg caa ccc gat ggc ggc aag acc gag aac tgc gcg gtc 481
Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala Val
              145              150              155

ctg tca ggc gcg gcc aac ggc aag tgg ttc gac aag cgc tgc cgc gat 529
Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp
              160              165              170

cag ctg ccc tac atc tgc cag ttc ggg atc gtg gcg gccgc 570
Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
175              180              185

```

<210> 9
 <211> 186
 <212> PRT
 <213> Homo sapiens

<400> 9

```

Pro Ala Met Ala Glu Pro Pro Thr Gln Lys Pro Lys Lys Ile Val Asn
  1             5             10             15
Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu Leu Lys Ser
      20             25             30
Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys Glu Gln Gln
      35             40             45
Ala Leu Gln Thr Val Cys Leu Lys Gly Thr Lys Val His Met Lys Cys
      50             55             60
Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala Ser Glu Asp
      65             70             75             80
Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr Gly Ser Glu
      85             90             95
Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly Asn Glu Ala
      100            105            110
Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Thr Trp Val
      115            120            125
Asp Met Thr Gly Ala Arg Ile Ala Tyr Lys Asn Trp Glu Thr Glu Ile
      130            135            140
Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala Val Leu Ser
      145            150            155            160
Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp Gln Leu
      165            170            175
Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
      180            185

```

<210> 10
 <211> 438
 <212> DNA
 <213> Homo sapiens

<220>

<221> CDS

<222> (8)..(433)

<223> PhTN3 encoding insert

<400> 10

```

ggccccag ccg gcc atg gcc gcc ctg cag acg gtc tgc ctg aag ggg acc 49
      Pro Ala Met Ala Ala Leu Gln Thr Val Cys Leu Lys Gly Thr
      1             5             10
aag gtg cac atg aaa tgc ttt ctg gcc ttc acc cag acg aag acc ttc 97
Lys Val His Met Lys Cys Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe
      15             20             25             30

```


[illegible]

```
<210> 11
<211> 142
<212> PRT
<213> Homo sapiens
```

```

<400> 11
Pro Ala Met Ala Ala Leu Gln Thr Val Cys Leu Lys Gly Thr Lys Val
  1             5             10             15
His Met Lys Cys Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu
          20             25             30
Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln
          35             40             45
Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val
  50             55             60
Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu
  65             70             75             80
Gly Thr Trp Val Asp Met Thr Gly Ala Arg Ile Ala Tyr Lys Asn Trp
          85             90             95
Glu Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys
          100             105             110
Ala Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys
          115             120             125

```

Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
 130 135 140

<210> 12
 <211> 570
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (8)..(565)
 <223> Phtlec encoding insert

<400> 12
 ggcccag ccg gcc atg gcc gag cca cca acc cag aag ccc aag aag att 49
 Pro Ala Met Ala Glu Pro Pro Thr Gln Lys Pro Lys Lys Ile
 1 5 10

gta aat gcc aag aaa gat gtt gtg aac aca aag atg ttt gag gag ctc 97
 Val Asn Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu Leu
 15 20 25 30

aag agc cgt ctg gac acc ctg gcc cag gag gtg gcc ctg ctg aag gag 145
 Lys Ser Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys Glu
 35 40 45

cag cag gcc ctg cag acg gtc gtc ctg aag ggg acc aag gtg cac atg 193
 Gln Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val His Met
 50 55 60

aaa gtc ttt ctg gcc ttc acc cag acg aag acc ttc cac gag gcc agc 241
 Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala Ser
 65 70 75

gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc cct cag act ggc 289
 Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr Gly
 80 85 90

tcg gag aac gac gcc ctg tat gag tac ctg cgc cag agc gtg ggc aac 337
 Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly Asn
 95 100 105 110

gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg gcc gag ggc acc 385
 Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Thr
 115 120 125

tgg gtg gac atg acc ggt acc cgc atc gcc tac aag aac tgg gag act 433
 Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys Asn Trp Glu Thr
 130 135 140

gag atc acc gcg caa ccc gat ggc ggc aag acc gag aac tgc gcg gtc 481
 Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala Val
 145 150 155

ctg tca ggc gcg gcc aac ggc aag tgg ttc gac aag cgc tgc cgc gat 529
 Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp
 160 165 170

caa ttg ccc tac atc tgc cag ttc ggg atc gtg gcg gccgc 570
 Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
 175 180 185

<210> 13
 <211> 186
 <212> PRT
 <213> Homo sapiens

<400> 13
 Pro Ala Met Ala Glu Pro Pro Thr Gln Lys Pro Lys Lys Ile Val Asn
 1 5 10 15
 Ala Lys Lys Asp Val Val Asn Thr Lys Met Phe Glu Glu Leu Lys Ser
 20 25 30
 Arg Leu Asp Thr Leu Ala Gln Glu Val Ala Leu Leu Lys Glu Gln Gln
 35 40 45
 Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val His Met Lys Val
 50 55 60
 Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu Ala Ser Glu Asp
 65 70 75 80
 Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln Thr Gly Ser Glu
 85 90 95
 Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val Gly Asn Glu Ala
 100 105 110
 Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Thr Trp Val
 115 120 125
 Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys Asn Trp Glu Thr Glu Ile
 130 135 140
 Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys Ala Val Leu Ser
 145 150 155 160
 Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp Gln Leu
 165 170 175
 Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
 180 185

<210> 14
 <211> 438
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (8)..(433)
 <223> PhtCTLD encoding insert

<400> 14

```

ggccccag ccg gcc atg gcc gcc ctg cag acg gtc gtc ctg aag ggg acc 49
      Pro Ala Met Ala Ala Leu Gln Thr Val Val Leu Lys Gly Thr
        1                      5                      10

aag gtg cac atg aaa gtc ttt ctg gcc ttc acc cag acg aag acc ttc 97
Lys Val His Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe
  15                      20                      25                      30

cac gag gcc agc gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc 145
His Glu Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr
                      35                      40                      45

cct cag act ggc tcg gag aac gac gcc ctg tat gag tac ctg cgc cag 193
Pro Gln Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln
                      50                      55                      60

agc gtg ggc aac gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg 241
Ser Val Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
      65                      70                      75

gcc gag ggc acc tgg gtg gac atg acc ggt acc cgc atc gcc tac aag 289
Ala Glu Gly Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys
      80                      85                      90

aac tgg gag act gag atc acc gcg caa ccc gat ggc ggc aag acc gag 337
Asn Trp Glu Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu
      95                      100                      105                      110

aac tgc gcg gtc ctg tca ggc gcg gcc aac ggc aag tgg ttc gac aag 385
Asn Cys Ala Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
                      115                      120                      125

cgc tgc cgc gat caa ttg ccc tac atc tgc cag ttc ggg atc gtg gcg 433
Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
                      130                      135                      140

gccgc 438

```

<210> 15

<211> 142

<212> PRT

<213> Homo sapiens

<400> 15

```

Pro Ala Met Ala Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val
  1                      5                      10                      15

His Met Lys Val Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu
      20                      25                      30

Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln
      35                      40                      45

Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val
      50                      55                      60

Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu
      65                      70                      75                      80

```

Gly Thr Trp Val Asp Met Thr Gly Thr Arg Ile Ala Tyr Lys Asn Trp
 85 90 95

Glu Thr Glu Ile Thr Ala Gln Pro Asp Gly Gly Lys Thr Glu Asn Cys
 100 105 110

Ala Val Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys
 115 120 125

Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Gly Ile Val Ala
 130 135 140

<210> 16

<211> 555

<212> DNA

<213> Mus musculus

<220>

<223> EcoRI to HindIII insert containing mtlec encoding
part

<400> 16

ggaattcgag tcacccactc ccaaggccaa gaaggctgca aatgccaaaga aagatttggt 60
 gagctcaaag atgtcgagga gctcaagaac aggatggatg tcttgGCCca ggaggtggcc 120
 ctgctgaagg agaagcaggc cttacagact gtggtcctga agggcaccAA ggtgaacttg 180
 aaggtcctcc tggccttcac ccaaccgaag accttccatg aggcgagcga ggactgcatc 240
 tcgcaagggg gcacgctggg caccocgcag tcagagctag agaacgaggc gctgttcgag 300
 tacgcgcgcc acagcgtggg caacgatgCG gagatctggc tgggcctcaa cgacatggcc 360
 gcggaaggcg cctgggtgga catgaccggT accctcctgg cctacaagaa ctgggagacg 420
 gagatcacga cgcaaccCGa cggcggcaaa gccgagaact gcgccgccct gtctggcgca 480
 gccaacggca agtggttcga caagcgatgc cgcgatcaat tgccctacat ctgccagttt 540
 gccattgtga agctt 555

<210> 17

<211> 77

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 17

cggaattcga gtcacccact cccaaggcca agaaggctgc aaatgccaaag aaagatttgg 60
 tgagctcaaa gatgttc 77

<210> 18

<211> 94

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 18

gcggatccag gcctgcttct ccttcagcag ggccacctcc tgggccagga catccatcct 60
 gttcttgagc tcctcgaaca tctttgagct cacc 94

<210> 19
<211> 97
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 19
gcaggcctta cagactgtgt gcctgaaggg caccaagggtg aacttgaagt gcctcctggc 60
cttcacccaa ccgaagacct tccatgaggc gagcgag 97

<210> 20
<211> 93
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 20
ccgcatgctt cgaacagcgc ctggttctct agctctgact gcgggggtgcc cagcgtgccc 60
ccttgcgaga tgcagtcctc gctcgctca tgg 93

<210> 21
<211> 61
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 21
ggttcgaata cgcgcgccac agcgtgggca acgatgcgga gatctaaatg ctcccaattg 60
c 61

<210> 22
<211> 55
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 22
ccaagcttca caatggcaaa ctggcagatg tagggcaatt gggagcattt agatc 55

<210> 23
<211> 86
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 23

cggagatctg gctgggcctc aacgacatgg ccgcggaagg cgctgggtg gacatgaccg 60
gtaccctcct ggcctacaag aactgg 86

<210> 24

<211> 130

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 24

gggcaattga tcgcggcatc gcttgtcgaa cctcttgccg ttggtgcgc cagacagggc 60
ggcgcagttc tcggctttgc cgccgtcggg ttgcgtcgtg atctccgtct cccagttctt 120
gtaggccagg 130

<210> 25

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 25

ctgggatcca tccagggtcg cgagtcaccc actcccaagg 40

<210> 26

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 26

ccgaagctta cacaatggca aactggc 27

<210> 27

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 27

ctgggatcca tccagggtcg cgccttacag actgtggtc 39

<210> 28
 <211> 568
 <212> DNA
 <213> Mus musculus

 <220>
 <221> CDS
 <222> (1)..(561)
 <223> FX-mtlec encoding insert

 <400> 28
 gga tcc atc cag ggt cgc gag tca ccc act ccc aag gcc aag aag gct 48
 Gly Ser Ile Gln Gly Arg Glu Ser Pro Thr Pro Lys Ala Lys Lys Ala
 1 5 10 15
 gca aat gcc aag aaa gat ttg gtg agc tca aag atg ttc gag gag ctc 96
 Ala Asn Ala Lys Lys Asp Leu Val Ser Ser Lys Met Phe Glu Glu Leu
 20 25 30
 aag aac agg atg gat gtc ctg gcc cag gag gtg gcc ctg ctg aag gag 144
 Lys Asn Arg Met Asp Val Leu Ala Gln Glu Val Ala Leu Leu Lys Glu
 35 40 45
 aag cag gcc tta cag act gtg gtc ctg aag ggc acc aag gtg aac ttg 192
 Lys Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val Asn Leu
 50 55 60
 aag gtc ctc ctg gcc ttc acc caa ccg aag acc ttc cat gag gcg agc 240
 Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe His Glu Ala Ser
 65 70 75 80
 gag gac tgc atc tcg caa ggg ggc acg ctg ggc acc ccg cag tca gag 288
 Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr Pro Gln Ser Glu
 85 90 95
 cta gag aac gag gcg ctg ttc gag tac gcg cgc cac agc gtg ggc aac 336
 Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His Ser Val Gly Asn
 100 105 110
 gat gcg gag atc tgg ctg ggc ctc aac gac atg gcc gcg gaa ggc gcc 384
 Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Ala
 115 120 125
 tgg gtg gac atg acc ggt acc ctc ctg gcc tac aag aac tgg gag acg 432
 Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys Asn Trp Glu Thr
 130 135 140
 gag atc acg acg caa ccc gac ggc ggc aaa gcc gag aac tgc gcc gcc 480
 Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu Asn Cys Ala Ala
 145 150 155 160
 ctg tct ggc gca gcc aac ggc aag tgg ttc gac aag cga tgc cgc gat 528
 Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp
 165 170 175
 caa ttg ccc tac atc tgc cag ttt gcc att gtg taagctt 568
 Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val
 180 185

<210> 29

<211> 187

<212> PRT

<213> Mus musculus

<400> 29

Gly Ser Ile Gln Gly Arg Glu Ser Pro Thr Pro Lys Ala Lys Lys Ala
 1 5 10 15

Ala Asn Ala Lys Lys Asp Leu Val Ser Ser Lys Met Phe Glu Glu Leu
 20 25 30

Lys Asn Arg Met Asp Val Leu Ala Gln Glu Val Ala Leu Leu Lys Glu
 35 40 45

Lys Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val Asn Leu
 50 55 60

Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe His Glu Ala Ser
 65 70 75 80

Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr Pro Gln Ser Glu
 85 90 95

Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His Ser Val Gly Asn
 100 105 110

Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Ala
 115 120 125

Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys Asn Trp Glu Thr
 130 135 140

Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu Asn Cys Ala Ala
 145 150 155 160

Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp
 165 170 175

Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val
 180 185

<210> 30

<211> 436

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1)..(429)

<223> FX-mtCTLD encoding insert

<400> 30

gga tcc atc cag ggt cgc gcc tta cag act gtg gtc ctg aag ggc acc 48
 Gly Ser Ile Gln Gly Arg Ala Leu Gln Thr Val Val Leu Lys Gly Thr
 1 5 10 15

aag gtg aac ttg aag gtc ctc ctg gcc ttc acc caa ccg aag acc ttc 96
 Lys Val Asn Leu Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe
 20 25 30

cat gag gcg agc gag gac tgc atc tcg caa ggg ggc acg ctg ggc acc 144
 His Glu Ala Ser Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr
 35 40 45

ccg cag tca gag cta gag aac gag gcg ctg ttc gag tac gcg cgc cac 192
 Pro Gln Ser Glu Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His
 50 55 60

agc gtg ggc aac gat gcg gag atc tgg ctg ggc ctc aac gac atg gcc 240
 Ser Val Gly Asn Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
 65 70 75 80

gcg gaa ggc gcc tgg gtg gac atg acc ggt acc ctc ctg gcc tac aag 288
 Ala Glu Gly Ala Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys
 85 90 95

aac tgg gag acg gag atc acg acg caa ccc gac ggc ggc aaa gcc gag 336
 Asn Trp Glu Thr Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu
 100 105 110

aac tgc gcc gcc ctg tct ggc gca gcc aac ggc aag tgg ttc gac aag 384
 Asn Cys Ala Ala Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
 115 120 125

cga tgc cgc gat caa ttg ccc tac atc tgc cag ttt gcc att gtg 429
 Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val
 130 135 140

taagctt 436

<210> 31

<211> 143

<212> PRT

<213> Mus musculus

<400> 31

Gly Ser Ile Gln Gly Arg Ala Leu Gln Thr Val Val Leu Lys Gly Thr
 1 5 10 15

Lys Val Asn Leu Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe
 20 25 30

His Glu Ala Ser Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr
 35 40 45

Pro Gln Ser Glu Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His
 50 55 60

Ser Val Gly Asn Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
 65 70 75 80

Ala Glu Gly Ala Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys
 85 90 95

Asn Trp Glu Thr Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu
 100 105 110

Asn Cys Ala Ala Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
 115 120 125

Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val
 130 135 140

<210> 32
 <211> 47
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 32
 cggctgagcg gccagccgg ccatggccga gtcacccact cccaagg 47

<210> 33
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 33
 cctgcggccg ccacgatccc gaactgg 27

<210> 34
 <211> 46
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 34
 cggctgagcg gccagccgg ccatggccgc cttacagact gtggtc 46

<210> 35
 <211> 570
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (8)..(565)
 <223> Pmtlec encoding insert

<400> 35
 ggccccag ccg gcc atg gcc gag tca ccc act ccc aag gcc aag aag gct 49
 Pro Ala Met Ala Glu Ser Pro Thr Pro Lys Ala Lys Lys Ala
 1 5 10

gca aat gcc aag aaa gat ttg gtg agc tca aag atg ttc gag gag ctc 97
 Ala Asn Ala Lys Lys Asp Leu Val Ser Ser Lys Met Phe Glu Glu Leu
 15 20 25 30

aag aac agg atg gat gtc ctg gcc cag gag gtg gcc ctg ctg aag gag 145
Lys Asn Arg Met Asp Val Leu Ala Gln Glu Val Ala Leu Leu Lys Glu
35 40 45

aag cag gcc tta cag act gtg gtc ctg aag ggc acc aag gtg aac ttg 193
Lys Gln Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val Asn Leu
50 55 60

aag gtc ctc ctg gcc ttc acc caa ccg aag acc ttc cat gag gcg agc 241
Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe His Glu Ala Ser
65 70 75

gag gac tgc atc tcg caa ggg ggc acg ctg ggc acc ccg cag tca gag 289
Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr Pro Gln Ser Glu
80 85 90

cta gag aac gag gcg ctg ttc gag tac gcg cgc cac agc gtg ggc aac 337
Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His Ser Val Gly Asn
95 100 105 110

gat gcg gag atc tgg ctg ggc ctc aac gac atg gcc gcg gaa ggc gcc 385
Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Ala
115 120 125

tgg gtg gac atg acc ggt acc ctc ctg gcc tac aag aac tgg gag acg 433
Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys Asn Trp Glu Thr
130 135 140

gag atc acg acg caa ccc gac ggc ggc aaa gcc gag aac tgc gcc gcc 481
Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu Asn Cys Ala Ala
145 150 155

ctg tct ggc gca gcc aac ggc aag tgg ttc gac aag cga tgc cgc gat 529
Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp
160 165 170

caa ttg ccc tac atc tgc cag ttt gcc att gtg gcg gccgc 570
Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val Ala
175 180 185

<210> 36
<211> 186
<212> PRT
<213> Mus musculus

<400> 36
Pro Ala Met Ala Glu Ser Pro Thr Pro Lys Ala Lys Lys Ala Ala Asn
1 5 10 15
Ala Lys Lys Asp Leu Val Ser Ser Lys Met Phe Glu Glu Leu Lys Asn
20 25 30
Arg Met Asp Val Leu Ala Gln Glu Val Ala Leu Leu Lys Glu Lys Gln
35 40 45
Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val Asn Leu Lys Val
50 55 60
Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe His Glu Ala Ser Glu Asp
65 70 75 80

Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr Pro Gln Ser Glu Leu Glu
85 90 95

Asn Glu Ala Leu Phe Glu Tyr Ala Arg His Ser Val Gly Asn Asp Ala
100 105 110

Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu Gly Ala Trp Val
115 120 125

Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys Asn Trp Glu Thr Glu Ile
130 135 140

Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu Asn Cys Ala Ala Leu Ser
145 150 155 160

Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys Arg Asp Gln Leu
165 170 175

Pro Tyr Ile Cys Gln Phe Ala Ile Val Ala
180 185

<210> 37

<211> 438

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (8)..(433)

<223> PmtCTLD encoding insert

<400> 37

ggcccag ccg gcc atg gcc gcc tta cag act gtg gtc ctg aag ggc acc 49
Pro Ala Met Ala Ala Leu Gln Thr Val Val Leu Lys Gly Thr
1 5 10

aag gtg aac ttg aag gtc ctc ctg gcc ttc acc caa ccg aag acc ttc 97
Lys Val Asn Leu Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe
15 20 25 30

cat gag gcg agc gag gac tgc atc tcg caa ggg ggc acg ctg ggc acc 145
His Glu Ala Ser Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr
35 40 45

ccg cag tca gag cta gag aac gag gcg ctg ttc gag tac gcg cgc cac 193
Pro Gln Ser Glu Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His
50 55 60

agc gtg ggc aac gat gcg gag atc tgg ctg ggc ctc aac gac atg gcc 241
Ser Val Gly Asn Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
65 70 75

gcg gaa ggc gcc tgg gtg gac atg acc ggt acc ctc ctg gcc tac aag 289
Ala Glu Gly Ala Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys
80 85 90

aac tgg gag acg gag atc acg acg caa ccc gac ggc ggc aaa gcc gag 337
Asn Trp Glu Thr Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu
95 100 105 110

aac tgc gcc gcc ctg tct ggc gca gcc aac ggc aag tgg ttc gac aag 385
 Asn Cys Ala Ala Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys
 115 120 125

cga tgc cgc gat caa ttg ccc tac atc tgc cag ttt gcc att gtg gcg 433
 Arg Cys Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val Ala
 130 135 140

gccgc 438

<210> 38
 <211> 142
 <212> PRT
 <213> Mus musculus

<400> 38
 Pro Ala Met Ala Ala Leu Gln Thr Val Val Leu Lys Gly Thr Lys Val
 1 5 10 15

Asn Leu Lys Val Leu Leu Ala Phe Thr Gln Pro Lys Thr Phe His Glu
 20 25 30

Ala Ser Glu Asp Cys Ile Ser Gln Gly Gly Thr Leu Gly Thr Pro Gln
 35 40 45

Ser Glu Leu Glu Asn Glu Ala Leu Phe Glu Tyr Ala Arg His Ser Val
 50 55 60

Gly Asn Asp Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu
 65 70 75 80

Gly Ala Trp Val Asp Met Thr Gly Thr Leu Leu Ala Tyr Lys Asn Trp
 85 90 95

Glu Thr Glu Ile Thr Thr Gln Pro Asp Gly Gly Lys Ala Glu Asn Cys
 100 105 110

Ala Ala Leu Ser Gly Ala Ala Asn Gly Lys Trp Phe Asp Lys Arg Cys
 115 120 125

Arg Asp Gln Leu Pro Tyr Ile Cys Gln Phe Ala Ile Val Ala
 130 135 140

<210> 39
 <211> 116
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 oligonucleotide

<400> 39
 cgctacaag aactggnsn nsnnnsnsn snnscaacc gatnsnsn nsnnsgagaa 60
 ctgcgcggtc ctgtcaggcg cggccaacgg caagtggns gacaagcgct gccgcg 116

<210> 40
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 40
gaccggtacc cgcacgcct acaagaactg g 31

<210> 41
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 41
gtagggaat tgatcgaggc agcgcttgc 30

<210> 42
<211> 94
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 42
gctgggcctc aacgacnnsn nsnsgagnn snnstgggtg gacatgaccg gtacccgcat 60
cgcctacaag aactgggaga ctgagatcac cgcg 94

<210> 43
<211> 102
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 43
cgcggcagcg cttgtcgaac cacttgccgt tggccgcgcc tgacaggacc gcgcagttct 60
csnnsnnsnn snnatcgggt tgcgcggtga tctcagtctc cc 102

<210> 44
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 44

cgaggccgag atctggctgg gcctcaacga c

31

<210> 45

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 45

gggcaacgag gccgagatct ggctgggcct c

31

<210> 46

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 46

cctgaccctg cagcgcttg

19

<210> 47

<211> 81

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 47

cgagatctgg ctgggcctca acgacnnsnn snnsnnsnns nnsaggaggca cctgggtgga 60
catgaccggt accgcatcg c 81

<210> 48

<211> 78

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 48

cgagatctgg ctgggcctca acgacnnsnn snnsnnsnns gagggcacct gggtggacat 60
gaccggtacc cgcacgc 78

<210> 49
<211> 94
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 49
gctgggcctc aacgacnnsn nsnnsgagnn snnstgggtg gacatgaccg gtaccgcat 60
cgcctacaag aactgggaga ctgagatcac cgcg 94

<210> 50
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 50
gcgatgcggg taccggtc 18

<210> 51
<211> 89
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 51
gcatagccta caagaactgg gagactgaga tcaccgogca acccgatggc ggcnnnsnnsn 60
nsnnnsnnsn sgagaactgc gcggtcctg 89

<210> 52
<211> 86
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 52
gcatagccta caagaactgg gagactgaga tcaccgogca acccgatggc ggcnnnsnnsn 60
nsnnnsnsga gaactgcgcg gtcctg 86

<210> 53
<211> 34
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 53

catgaccggt acccgcatcg cctacaagaa ctgg

34

<210> 54

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 54

cctgaccctg cagcgcttgt cgaaccactt gccgttggcc gcgcctgaca ggaccgcgca 60
gttctc 66

<210> 55

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 55

ggtacctaag tgacgatatc ctgacctaac tgcagggatc aattg

45

<210> 56

<211> 343

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (8)..(274)

<223> Human PhtCPB insert

<400> 56

ggcccag ccg gcc atg gcc gcc ctc cag acg gtc tgc ctg aag ggg acc 49
Pro Ala Met Ala Ala Leu Gln Thr Val Cys Leu Lys Gly Thr
1 5 10

aag gtg cac atg aaa tgc ttt ctg gcc ttc acc cag acg aag acc ttc 97
Lys Val His Met Lys Cys Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe
15 20 25 30

cac gag gcc agc gag gac tgc atc tcg cgc ggg ggc acc ctg agc acc 145
His Glu Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr
35 40 45

cct cag act ggc tcg gag aac gac gcc ctg tat gag tac ctg cgc cag 193
Pro Gln Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln
50 55 60

25

agc gtg ggc aac gag gcc gag atc tgg ctg ggc ctc aac gac atg gcg 241
Ser Val Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala
65 70 75

gcc gag ggc acc tgg gtg gac atg acc ggt acc taagtgcga tatectgacc 294
Ala Glu Gly Thr Trp Val Asp Met Thr Gly Thr
80 85

taactgcagg gatcaattgc cctacatctg ccagttcggg atcgtgtag 343

```
<210> 57
<211> 89
<212> PRT
<213> Homo sapiens
```

```

<400> 57
Pro Ala Met Ala Ala Leu Gln Thr Val Cys Leu Lys Gly Thr Lys Val
  1             5             10             15
His Met Lys Cys Phe Leu Ala Phe Thr Gln Thr Lys Thr Phe His Glu
             20             25             30
Ala Ser Glu Asp Cys Ile Ser Arg Gly Gly Thr Leu Ser Thr Pro Gln
             35             40             45
Thr Gly Ser Glu Asn Asp Ala Leu Tyr Glu Tyr Leu Arg Gln Ser Val
  50             55             60
Gly Asn Glu Ala Glu Ile Trp Leu Gly Leu Asn Asp Met Ala Ala Glu
  65             70             75             80
Gly Thr Trp Val Asp Met Thr Gly Thr
             85

```

```
<210> 58
<211> 405
<212> DNA
<213> Rattus rattus
```

```
<220>
<221> CDS
<222> (8)..(400)
<223> Rat PrMBP insert
```

<400> 58																
ggcccag	ccg	gcc	atg	gcc	aac	aag	ttg	cat	gcc	ttc	tcc	atg	ggg	aaa		49
	Pro	Ala	Met	Ala	Asn	Lys	Leu	His	Ala	Phe	Ser	Met	Gly	Lys		
	1				5					10						
aag	tct	ggg	aag	aag	ttc	ttt	gtg	acc	aac	cat	gaa	agg	atg	ccc	ttt	97
Lys	Ser	Gly	Lys	Lys	Phe	Phe	Val	Thr	Asn	His	Glu	Arg	Met	Pro	Phe	
15					20					25					30	
tcc	aaa	gtc	aag	gcc	ctg	tgc	tca	gag	ctc	cga	ggc	act	gtg	gct	atc	145
Ser	Lys	Val	Lys	Ala	Leu	Cys	Ser	Glu	Leu	Arg	Gly	Thr	Val	Ala	Ile	
				35				40					45			

ccc aag aat gct gag gag aac aag gcc atc caa gaa gtg gct aaa acc 193
 Pro Lys Asn Ala Glu Glu Asn Lys Ala Ile Gln Glu Val Ala Lys Thr
 50 55 60

tct gcc ttc cta ggc atc acg gac gag gtg act gaa ggc caa ttc atg 241
 Ser Ala Phe Leu Gly Ile Thr Asp Glu Val Thr Glu Gly Gln Phe Met
 65 70 75

tat gtg aca ggg ggg agg ctc acc tac agc aac tgg aaa aag gat gag 289
 Tyr Val Thr Gly Gly Arg Leu Thr Tyr Ser Asn Trp Lys Lys Asp Glu
 80 85 90

ccc aat gac cat ggc tct ggg gaa gac tgt gtc act ata gta gac aac 337
 Pro Asn Asp His Gly Ser Gly Glu Asp Cys Val Thr Ile Val Asp Asn
 95 100 105 110

ggt ctg tgg aat gac atc tcc tgc caa gct tcc cac acg gct gtc tgc 385
 Gly Leu Trp Asn Asp Ile Ser Cys Gln Ala Ser His Thr Ala Val Cys
 115 120 125

gag ttc cca gcc gcg gccgc 405
 Glu Phe Pro Ala Ala
 130

<210> 59
 <211> 131
 <212> PRT
 <213> Rattus rattus

<400> 59
 Pro Ala Met Ala Asn Lys Leu His Ala Phe Ser Met Gly Lys Lys Ser
 1 5 10 15

Gly Lys Lys Phe Phe Val Thr Asn His Glu Arg Met Pro Phe Ser Lys
 20 25 30

Val Lys Ala Leu Cys Ser Glu Leu Arg Gly Thr Val Ala Ile Pro Lys
 35 40 45

Asn Ala Glu Glu Asn Lys Ala Ile Gln Glu Val Ala Lys Thr Ser Ala
 50 55 60

Phe Leu Gly Ile Thr Asp Glu Val Thr Glu Gly Gln Phe Met Tyr Val
 65 70 75 80

Thr Gly Gly Arg Leu Thr Tyr Ser Asn Trp Lys Lys Asp Glu Pro Asn
 85 90 95

Asp His Gly Ser Gly Glu Asp Cys Val Thr Ile Val Asp Asn Gly Leu
 100 105 110

Trp Asn Asp Ile Ser Cys Gln Ala Ser His Thr Ala Val Cys Glu Phe
 115 120 125

Pro Ala Ala
 130

<210> 60
 <211> 408
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (8)..(403)
 <223> Human PhSP-D insert

<400> 60
 ggcccag ccg gcc atg gcc aag aaa gtt gag ctc ttc cca aat ggc caa 49
 Pro Ala Met Ala Lys Lys Val Glu Leu Phe Pro Asn Gly Gln
 1 5 10

 agt gtg ggg gag aag att ttc aag aca gca ggc ttt gta aaa cca ttt 97
 Ser Val Gly Glu Lys Ile Phe Lys Thr Ala Gly Phe Val Lys Pro Phe
 15 20 25 30

 acg gag gca cag ctg ctg tgc aca cag gct ggt gga cag ttg gcc tct 145
 Thr Glu Ala Gln Leu Leu Cys Thr Gln Ala Gly Gly Gln Leu Ala Ser
 35 40 45

 cca cgc tct gcc gct gag aat gcc gcc ttg caa cag ctg gtc gta gct 193
 Pro Arg Ser Ala Ala Glu Asn Ala Ala Leu Gln Gln Leu Val Val Ala
 50 55 60

 aag aac gag gct gct ttc ctg agc atg act gat tcc aag aca gag ggc 241
 Lys Asn Glu Ala Ala Phe Leu Ser Met Thr Asp Ser Lys Thr Glu Gly
 65 70 75

 aag ttc acc tac ccc aca gga gag tcc ctg gtc tat tcc aac tgg gcc 289
 Lys Phe Thr Tyr Pro Thr Gly Glu Ser Leu Val Tyr Ser Asn Trp Ala
 80 85 90

 cca ggg gag ccc aac gat gat ggc ggg tca gag gac tgt gtg gag atc 337
 Pro Gly Glu Pro Asn Asp Asp Gly Gly Ser Glu Asp Cys Val Glu Ile
 95 100 105 110

 ttc acc aat ggc aag tgg aat gac agg gct tgt gga gaa aag cgt ctt 385
 Phe Thr Asn Gly Lys Trp Asn Asp Arg Ala Cys Gly Glu Lys Arg Leu
 115 120 125

 gtg gtc tgc gag ttc gcg gccgc 408
 Val Val Cys Glu Phe Ala
 130

<210> 61
 <211> 132
 <212> PRT
 <213> Homo sapiens

<400> 61
 Pro Ala Met Ala Lys Lys Val Glu Leu Phe Pro Asn Gly Gln Ser Val
 1 5 10 15

 Gly Glu Lys Ile Phe Lys Thr Ala Gly Phe Val Lys Pro Phe Thr Glu
 20 25 30

Ala Gln Leu Leu Cys Thr Gln Ala Gly Gly Gln Leu Ala Ser Pro Arg
 35 40 45

Ser Ala Ala Glu Asn Ala Ala Leu Gln Gln Leu Val Val Ala Lys Asn
 50 55 60

Glu Ala Ala Phe Leu Ser Met Thr Asp Ser Lys Thr Glu Gly Lys Phe
 65 70 75 80

Thr Tyr Pro Thr Gly Glu Ser Leu Val Tyr Ser Asn Trp Ala Pro Gly
 85 90 95

Glu Pro Asn Asp Asp Gly Gly Ser Glu Asp Cys Val Glu Ile Phe Thr
 100 105 110

Asn Gly Lys Trp Asn Asp Arg Ala Cys Gly Glu Lys Arg Leu Val Val
 115 120 125

Cys Glu Phe Ala
 130

<210> 62
 <211> 49
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 oligonucleotide

<400> 62
 cggctgagcg gccagccgg ccatggccaa caagttgcat gccttctcc 49

<210> 63
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 oligonucleotide

<400> 63
 gcactcctgc ggccgcggct gggaactcgc agac 34

<210> 64
 <211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 oligonucleotide

<400> 64
 cggctgagcg gccagccgg ccatggccaa gaaagttgag ctcttccc 48

<210> 65
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 65
gcactcctgc ggccgcgaac tcgcagacca caagac

36

<210> 66
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 66
gccaccggtg acgtagatga attggccttc snnnsnnsnns nnsnngtccg tgatgcctag 60
gaagg 65

<210> 67
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 67
gccaccggtg acgtagatga attggccttc snnnsnnsnns nnsnnsnngt ccgtgatgcc 60
taggaagg 68

<210> 68
<211> 62
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 68
gccaccggtg acgtagatga asnnnsnnsnn snnnsnnsnns nncgtgatgc ctaggaaggc 60
ag 62

<210> 69
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 69
ccagttgctg tatttcaggc tgccaccggt gacgtagatg

40

<210> 70
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 70
gcctgaaata cagcaactgg aagaaagacg aacc

34

<210> 71
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 71
ctggaagaaa gacgaaccga atgaccatgg cnnsnnsnns nnsnnsagaag actgtgtcac 60
tatagtag 68

<210> 72
<211> 71
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 72
ctggaagaaa gacgaaccga atgaccatgg cnnsnnsnns nnsnnsnsg aagactgtgt 60
cactatagta g 71

<210> 73
<211> 59
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 73
ctggaagaaa gacgaaccga atnnsnnsnn snnsnnsnsgaa gactgtgtca ctatagtag 59

<210> 74
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 74
cggctgagcg gccacgc

17

<210> 75
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 75
gcactcctgc ggccgcg

17

<210> 76
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 76
ctcacgggtc ggatacgtga acttgccctc tgtsnnsnns nnsnnsnnat cagtcagtgc 60
caggaaagc 69

<210> 77
<211> 72
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 77
ctcacgggtc ggatacgtga acttgccctc tgtsnnsnns nnsnnsnnsn natcagtcag 60
gctcaggaaa gc 72

<210> 78
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:

oligonucleotide

<400> 78

ctcaccgggtc ggatacgtga asnnsnnsnn snnsnnsnns nnagtcatgc tcaggaaagc 60

<210> 79

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 79

cagttggaat agaccaggga ctcaccgggtc ggatacgtg

39

<210> 80

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 80

gggccccagg ggagcccaac gatgatggcn nsnnsnnsnn snnsgaggac tgtgtggaga 60
tcttc 65

<210> 81

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 81

gggccccagg ggagcccaac gatgatggcn nsnnsnnsnn snnsnnsnsgag gactgtgtgg 60
agatcttc 68

<210> 82

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 82

gggccccagg ggagcccaac gatgatggcn nsnnsnnsnn snnsnnsnsgag gactgtgtgg 60
agatcttc 68

<210> 83
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 83
gggccccagg ggagcccaac nnsnnsnnsn nsnnsgagga ctgtgtggag atcttc 56

<210> 84
<211> 77
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 84
gcatcgccta caagaactgg nnsnnsnnsn nsnnnsnsca acccgatggc ggcaagaccg 60
agaactgcgc ggtcctg 77

<210> 85
<211> 83
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 85
gcatcgccta caagaactgg gagnnsnnsn nsnnnsnnsn sgcgcaaccc gatggcgga 60
agaccgagaa ctgcgcggtc ctg 83

<210> 86
<211> 80
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide

<400> 86
gcatcgccta caagaactgg gagnnsnnsn nsnnnsnsgc gcaaccgat ggcggaaga 60
ccgagaactg cgcggtcctg 80

<210> 87
<211> 75
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 87

gtagggcaat tgatcgctgc agcgcttgtc gaaccasnns nnsnnsnnsn nsnnsnncag 60
gaccgcgcag ttctc 75

<210> 88

<211> 84

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 88

gtagggcaat tgatcgctgc agcgcttgtc gaaccacttg ccsnnsnnsn nsnnsnnsnn 60
gcctgacagg accgcgcagt tctc 84

<210> 89

<211> 81

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 89

gtagggcaat tgatcgctgc agcgcttgtc gaaccacttg ccsnnsnnsn nsnnsnngcc 60
tgacaggacc gcgcagttct c 81

<210> 90

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 90

gtagggcaat tgatcgctgc 20

<210> 91

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide

<400> 91

catgaccggt accgcacatcg cctacaagaa ctgg

**(19) World Intellectual Property Organization
International Bureau**



(43) International Publication Date
20 June 2002 (20.06.2002)

PCT

(10) International Publication Number
WO 02/048189 A3

(51) International Patent Classification⁷: C12N 15/12,
15/10, 15/62, C07K 14/47

(21) International Application Number: PCT/DK01/00825

(22) International Filing Date:
13 December 2001 (13.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2000 01872 13 December 2000 (13.12.2000) DK
60/272,098 28 February 2001 (28.02.2001) US

(71) Applicant (for all designated States except US): BOREAL PHARMA A/S [DK/DK]; Science Park Aarhus, Gustavs Wieds Vej 10, DK-8000 Aarhus C (DK).

(72) Inventors; and

(75) **Inventors/Applicants (for US only): ETZERODT, Michael [DK/DK]; Teglthøj 18, DK-8382 Hinnerup (DK). HOLTET, Thor, Las [DK/DK]; Roskildevej**

104, DK-3600 Frederikssund (DK). **GRAVERSEN, Niels, Jonas, Heilskov** [DK/DK]; Brendstrupvej 58, DK-8200 Århus N (DK). **THØGERSEN, Hans, Christian** [DK/DK]; Ristrupvej 41, DK-8381 Mundelstrup (DK).


(74) Agent: **HOFMAN-BANG ZACCO A/S**; Hans
Bekkevoolds Allé 7, DK-2900 Hellerup (DK).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: COMBINATORIAL LIBRARIES OF PROTEINS HAVING THE SCAFFOLD STRUCTURE OF C-TYPE LECTIN-LIKE DOMAINS



(57) Abstract: A novel family of protein libraries comprising CTLDS (C-type Lectin-Like Domains) in which internal polypeptide loop-regions lining the ligand binding sites in CTLDs have been replaced with ensembles of completely or partially randomised polypeptide segments. Tetranectin CTLDs were chosen as framework for the preferred embodiment of the invention; and versatile phagemid vectors useful in the generation and manipulation of human and murine tetranectin CTLD libraries are disclosed as part of this invention. Tetranectin CTLDs in monomeric as well as in trimeric form are efficiently displayed as gene III fusions in fully functional form by the recombinant fd phage display vector. CTLD derivatives with affinity for new ligands may readily be isolated from libraries of vectors displaying CTLDs, in which loop-regions have been randomised, using one or more rounds of enrichment by screening or selection followed by amplification of the enriched subpopulation in each round. The efficiency with which protein products containing CTLDs with new binding properties can be produced, e.g. by bacterial expression in *in vitro* refolding, in mono-, tri-, or multimeric formats provides important advantages in terms of simplicity, cost and efficiency of generation, production and diagnostic or therapeutic applications in comparison to recombinant antibody derivatives.

WO 02/048189 A3



Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:

9 January 2003

INTERNATIONAL SEARCH REPORT

national Application No

PCT/DK 01/00825

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/10 C12N15/62 C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CAB Data, SEQUENCE SEARCH, BIOSIS, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GRAVERSEN JONAS HEILSKOV ET AL: "Mutational analysis of affinity and selectivity of kringle-tetranectin interaction. Grafting novel kringle affinity onto the tetranectin lectin scaffold." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 275, no. 48, 1 December 2000 (2000-12-01), pages 37390-37396, XP002175478 ISSN: 0021-9258 cited in the application the whole document</p> <p style="text-align: center;">--- -/--</p>	<p>1-4,6,7, 12,13, 39,58</p>



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

14 August 2002

Date of mailing of the international search report

27/08/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-2016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

national Application No
PCT/DK 01/00825

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GRAVERSEN JONAS HEILSKOV ET AL: "The plasminogen binding site of the C-type lectin tetranectin is located in the carbohydrate recognition domain, and binding is sensitive to both calcium and lysine."</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 44, 30 October 1998 (1998-10-30), pages 29241-29246, XP002175479 ISSN: 0021-9258 cited in the application the whole document</p> <p style="text-align: center;">---</p>	<p>1-4,6,7, 12,13, 39,58</p>
X	<p>TORGENSEN DAWN ET AL: "Mechanism of ligand binding to E- and P-selectin analyzed using selectin/mannose-binding protein chimeras."</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 11, 13 March 1998 (1998-03-13), pages 6254-6261, XP002209822 ISSN: 0021-9258 cited in the application the whole document</p> <p style="text-align: center;">---</p>	<p>1-5</p>
X	<p>IOBST SUSANNE T ET AL: "Binding of sugar ligands to Ca-2+-dependent animal lectins: I. Analysis of mannose binding by site-directed mutagenesis and NMR."</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 22, 1994, pages 15505-15511, XP002209823 ISSN: 0021-9258 cited in the application the whole document</p> <p style="text-align: center;">---</p>	<p>1-5</p>
X	<p>IOBST SUSANNE T ET AL: "Binding of sugar ligands to Ca-2+-dependent animal lectins: II. Generation of high-affinity galactose binding by site-directed mutagenesis."</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 22, 1994, pages 15512-15519, XP002209824 ISSN: 0021-9258 cited in the application the whole document</p> <p style="text-align: center;">---</p>	<p>1-5</p>
A	<p>WO 98 56906 A (NIELSEN BETTINA BRYDE ;THOEGENSEN HANS CHRISTIAN (DK); ETZERODT MI) 17 December 1998 (1998-12-17) cited in the application claims 1-55</p> <p style="text-align: center;">---</p>	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 01/00825

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 18227 A (DENZYME APS ;HOLTET THOR LAS (DK); ETZERODT MICHAEL (DK); THOGERSE) 18 August 1994 (1994-08-18) cited in the application claims 1-66; figures 20,21; example 9 ---	
A	DOI N ET AL: "SCREENING OF CONFORMATIONALLY CONSTRAINED RANDOM POLYPEPTIDE LIBRARIES DISPLAYED ON A PROTEIN SCAFFOLD" CMLS CELLULAR AND MOLECULAR LIFE SCIENCES, BIRKHAUSER VERLAG, BASEL, CH, vol. 54, no. 5, May 1998 (1998-05), pages 394-404, XP000891835 ISSN: 1420-682X the whole document ---	
A	DRICKAMER KURT: "C-type lectin-like domains." CURRENT OPINION IN STRUCTURAL BIOLOGY, vol. 9, no. 5, October 1999 (1999-10), pages 585-590, XP002175480 ISSN: 0959-440X cited in the application the whole document ---	
A	WO 99 24617 A (KAMB CARL ALEXANDER ;ABEDI MAJID (US); VENTANA GENETICS INC (US)) 20 May 1999 (1999-05-20) the whole document ---	
A	WO 00 20574 A (RIGEL PHARMACEUTICALS INC) 13 April 2000 (2000-04-13) the whole document ---	
A	EP 0 947 582 A (INNOGENETICS NV) 6 October 1999 (1999-10-06) the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/DK 01/00825

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9856906	A	17-12-1998	AU 736707 B2	02-08-2001
			AU 7906598 A	30-12-1998
			CA 2304254 A1	17-12-1998
			WO 9856906 A1	17-12-1998
			EP 1012280 A1	28-06-2000
WO 9418227	A	18-08-1994	AU 674568 B2	02-01-1997
			AU 6038094 A	29-08-1994
			CA 2155335 A1	18-08-1994
			WO 9418227 A2	18-08-1994
			EP 0686162 A1	13-12-1995
			FI 953705 A	03-08-1995
			JP 8506243 T	09-07-1996
			NO 952989 A	03-10-1995
			NZ 261571 A	24-03-1997
			US 5739281 A	14-04-1998
			US 5917018 A	29-06-1999
			AT 199392 T	15-03-2001
			AT 173740 T	15-12-1998
			AU 690171 B2	23-04-1998
			AU 1117095 A	19-06-1995
			AU 690528 B2	30-04-1998
			AU 5654894 A	04-07-1994
			AU 680685 B2	07-08-1997
			AU 7621494 A	10-04-1995
			CA 2169620 A1	30-03-1995
			CA 2177367 A1	08-06-1995
			DE 69329974 D1	05-04-2001
			DE 69329974 T2	19-07-2001
			DE 69414870 D1	07-01-1999
			DE 69414870 T2	12-05-1999
			DK 672142 T3	18-06-2001
			DK 720624 T3	09-08-1999
			EP 0672142 A1	20-09-1995
			EP 0720624 A1	10-07-1996
			EP 0731842 A1	18-09-1996
			ES 2126145 T3	16-03-1999
			WO 9508577 A1	30-03-1995
			WO 9515388 A1	08-06-1995
			JP 9503759 T	15-04-1997
			JP 8504100 T	07-05-1996
			JP 9506508 T	30-06-1997
			US 5837242 A	17-11-1998
WO 9924617	A	20-05-1999	US 6025485 A	15-02-2000
			AU 1520099 A	31-05-1999
			CA 2309543 A1	20-05-1999
			EP 1029081 A1	23-08-2000
			JP 2001522587 T	20-11-2001
			NO 20002375 A	30-06-2000
			WO 9924617 A1	20-05-1999
WO 0020574	A	13-04-2000	US 6180343 B1	30-01-2001
			AU 1516400 A	26-04-2000
			EP 1119617 A2	01-08-2001
			WO 0020574 A2	13-04-2000
			US 2001003650 A1	14-06-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PLT/DK 01/00825

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
EP 0947582	A	06-10-1999	EP 0947582 A1	06-10-1999